



## Exact solutions of distinct physical structures to the fractional potential Kadomtsev–Petviashvili equation

**Ahmet Bekir**

Eskisehir Osmangazi University, Art-Science Faculty, Department of Mathematics-Computer, Eskisehir-TURKEY

E-mail: abekir@ogu.edu.tr;

**Özkan Güner**

Dumlupinar University, School of Applied Sciences, Department of Management Information Systems, Kutahya-TURKEY

E-mail: ozkan.guner@dpu.edu.tr

---

**Abstract** In this paper, Exp-function and  $\left(\frac{G'}{G}\right)$ -expansion methods are presented to derive traveling wave solutions for a class of nonlinear space-time fractional differential equations. As a results, some new exact traveling wave solutions are obtained.

---

**Keywords.** Exact solution; modified Riemann-Liouville derivative; solitons; space-time fractional potential Kadomtsev-Petviashvili (pKP) equation.

**2010 Mathematics Subject Classification.** 26A33, 34A08, 35R11, 83C15.

### 1. INTRODUCTION

Recently, theory and applications of fractional differential equations (FDEs) has been the focus of many studies due to their frequent appearance in various applications in mathematics, physics, biology, engineering, signal processing, systems identification, control theory, finance and fractional dynamics, and has attracted much attention of more and more scholars. The fractional differential equations (FDEs) have been investigated by many researchers [1, 2, 3]. Recent investigations show that the dynamic of many physical processes is described accurately by using fractional differential equations containing different types of fractional derivatives. The most popular derivatives of fractional order are the Caputo derivative, the Riemann-Liouville derivative and Grünwald-Letnikov derivative. A few years ago, Jumarie presented a different definition of the fractional derivative being a little modification of the Riemann-Liouville derivative.

Since fractional differential equations are used to describe a large variety of physical phenomena, finding exact solutions to FDEs is an important subject and a hot topic. Many powerful and efficient methods have been proposed so far including the fractional  $\left(\frac{G'}{G}\right)$ -expansion method [4, 5, 6, 7], the fractional exp-function method [8, 9, 10], the fractional first integral method [12, 13], the fractional sub-equation method [14, 15, 16], the fractional functional variable method [17], the fractioanal

modified trial equation method [18, 19, 20], the fractional simplest equation method [21] and so on. Using these methods, solutions with various forms for given fractional differential equation have been established.

The organization of this paper is as follows. In section 2, we give the basic definitions and analysis of methods, then in section 3, we give applications. Some conclusions are given in last section.

## 2. BASIC DEFINITIONS AND ANALYSIS OF METHODS

The Jumarie’s modified Riemann–Liouville derivative [22] of order  $\alpha$  is defined by the following expression:

$$D_t^\alpha f(t) = \begin{cases} \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-\xi)^{-\alpha} (f(\xi) - f(0)) d\xi & , \quad 0 < \alpha < 1 \\ (f^{(n)}(t))^{(\alpha-n)} & , \quad n \leq \alpha < n+1, \quad n \geq 1. \end{cases} \quad (2.1)$$

Now, some important properties of the fractional modified Riemann–Liouville derivative were summarized [23]

$$D_t^\alpha t^r = \frac{\Gamma(1+r)}{\Gamma(1+r-\alpha)} t^{r-\alpha} \quad , \quad (2.2)$$

$$D_t^\alpha (f(t)g(t)) = g(t)D_t^\alpha f(t) + f(t)D_t^\alpha g(t), \quad (2.3)$$

$$D_t^\alpha f[g(t)] = f'_g[g(t)]D_t^\alpha g(t) = D_g^\alpha [g(t)](g'(t))^\alpha. \quad (2.4)$$

The above equations play an important role in fractional calculus in the following applications.

Firstly we consider the following general nonlinear FDE of the type

$$P(u, D_t^\alpha u, D_x^\beta u, D_t^\alpha D_t^\alpha u, D_t^\alpha D_x^\beta u, D_x^\beta D_x^\beta u, \dots) = 0, \quad 0 < \alpha, \beta \leq 1 \quad (2.5)$$

where  $u$  is an unknown function. Moreover,  $P$  is a polynomial of  $u$  and its partial fractional derivatives, in which the highest order derivatives and the nonlinear terms are involved.

Li and He [24, 25] proposed a fractional complex transform to convert fractional differential equations into ordinary differential equations (ODEs). So all analytical methods devoted to the advanced calculus can be easily applied to the fractional calculus. The traveling wave variable

$$u(x, t) = U(\xi), \quad (2.6)$$

$$\xi = \frac{kx^\beta}{\Gamma(1+\beta)} + \frac{\tau t^\alpha}{\Gamma(1+\alpha)}, \quad (2.7)$$



where  $\tau$  and  $k$  are nonzero arbitrary constants, we can rewrite Eq. (2.5) in the following nonlinear ODE;

$$Q(U, U', U'', U''', \dots) = 0, \quad (2.8)$$

where the prime denotes the derivation with respect to  $\xi$ .

According to exp-function method, which was developed by He and Wu [26], we assume that the wave solution can be expressed in the following form

$$U(\xi) = \frac{\sum_{n=-c}^d a_n \exp[n\xi]}{\sum_{m=-p}^q b_m \exp[m\xi]} \quad (2.9)$$

where  $p, q, c$  and  $d$  are positive integers which are known to be further determined,  $a_n$  and  $b_m$  are unknown constants. We can rewrite Eq. (2.9) in the following equivalent form.

$$U(\xi) = \frac{a_{-c} \exp[-c\xi] + \dots + a_d \exp[d\xi]}{b_{-p} \exp[-p\xi] + \dots + b_q \exp[q\xi]} \quad (2.10)$$

This equivalent formulation plays an important and fundamental part for finding the analytic solution of problems. To determine the value of  $c$  and  $p$ , we balance the linear term of highest order of equation Eq. (2.10) with the highest order nonlinear term. Similarly, to determine the value of  $d$  and  $q$ , we balance the linear term of lowest order of Eq. (2.10) with lowest order nonlinear term [27, 28].

Secondly suppose the solution of equation (2.8) can be expressed by a polynomial in  $\left(\frac{G'}{G}\right)$  as follows:

$$U(\xi) = \sum_{i=0}^m a_i \left(\frac{G'}{G}\right)^i, \quad a_m \neq 0, \quad (2.11)$$

where  $a_i$  ( $i = 0, 1, 2, \dots, m$ ) are constants, while  $G(\xi)$  satisfies the following second order linear ordinary differential equation

$$G''(\xi) + \lambda G'(\xi) + \mu G(\xi) = 0, \quad (2.12)$$

where  $\lambda$  and  $\mu$  are constants. Then the positive integer  $m$  can be determined by considering the homogeneous balance between the highest order derivatives and the nonlinear terms appearing in equation (2.8). By substituting equation (2.11) into equation (2.8) and using equation (2.12), we collect all terms with the same order of  $\left(\frac{G'}{G}\right)$ . Then by equating each coefficient of the resulting polynomial to zero, we obtain a set of algebraic equations for  $a_i$  ( $i = 0, 1, 2, \dots, m$ ),  $\lambda$ ,  $\mu$ ,  $\tau$  and  $k$ . By solving the equations system, substituting  $a_i$  ( $i = 0, 1, 2, \dots, m$ ),  $\lambda$ ,  $\mu$ ,  $\tau$ ,  $k$  and the general solutions of equation (2.12) into equation (2.11), we can get a variety of exact solutions of equation (2.5) [29, 30].



### 3. APPLICATIONS

We consider the space-time fractional potential Kadomtsev–Petviashvili (pKP) equation [31] in the form:

$$\frac{1}{4}D_x^{4\alpha}u + \frac{3}{2}D_x^\alpha u D_x^{2\alpha}u + \frac{3}{4}D_y^{2\alpha}u + D_t^\alpha(D_x^\alpha u) = 0, \quad t > 0, \quad 0 < \alpha \leq 1, \quad (3.1)$$

where  $\alpha$  is a parameter describing the order of the fractional space-time derivative. When  $\alpha = 1$  in Eq. (3.1) is the fractional differential equation reduces to the KP type equation [32-37].

For our purpose, we introduce the following transformations;

$$u(x, y, t) = U(\xi) \quad (3.2)$$

$$\xi = \frac{kx^\alpha}{\Gamma(1 + \alpha)} + \frac{ny^\alpha}{\Gamma(1 + \alpha)} + \frac{ct^\alpha}{\Gamma(1 + \alpha)}, \quad (3.3)$$

where  $k$ ,  $n$  and  $c$  are nonzero constants.

Substituting (3.3) into (3.1), reduces (3.1) into an ODE

$$\frac{k^4}{4}U'''' + \frac{3k^3}{2}U'U'' + \frac{3n^2}{4}U'' + ckU'' = 0, \quad (3.4)$$

where  $U'' = \frac{dU}{d\xi}$ .

Integrating equation (3.4) with respect to  $\xi$  yields

$$\frac{k^4}{4}U''' + \frac{3k^3}{4}(U')^2 + (\frac{3n^2}{4} + ck)U' + \xi_0 = 0, \quad (3.5)$$

where  $\xi_0$  is a constant of integration.

Firstly we begin the pKP equation to solve by using the exp-function method. We can determine values of  $d$  and  $q$  by balancing the order of  $U'''$  and  $(U')^2$  in Eq.(3.5), we get

$$U''' = \frac{d_1 \exp[(7q + d)\xi] + \dots}{d_2 \exp[8q\xi] + \dots}, \quad (3.6)$$

and

$$(U')^2 = \frac{d_3 \exp[2(d + q)\xi] + \dots}{d_4 \exp[4q\xi] + \dots}, \quad (3.7)$$

where  $d_i$  are determined coefficients only for simplicity. Balancing highest order of Exp-function in Eqs.(3.6) and (3.7) we obtain

$$7q + d = 2d + 6q, \quad (3.8)$$

which leads to the result

$$q = d. \quad (3.9)$$



In the same way, to determine values of  $c$  and  $p$ , we balance the linear term of the lowest order in Eq.(3.5),

$$U''' = \frac{\dots + c_1 \exp[-(7p + c)\xi]}{\dots + c_2 \exp[-8p\xi]}, \quad (3.10)$$

and

$$(U')^2 = \frac{\dots + c_3 \exp[-2(c + p)\xi]}{\dots + c_4 \exp[-4p\xi]}, \quad (3.11)$$

where  $c_i$  are determined coefficients only for simplicity. From (3.10) and (3.11), we have

$$-7p - c = -2c - 6p, \quad (3.12)$$

and this gives

$$p = c. \quad (3.13)$$

For simplicity, we set  $p = c = 1$  and  $q = d = 1$ , so Eq.(2.6) reduces to

$$U(\xi) = \frac{a_1 \exp(\xi) + a_0 + a_{-1} \exp(-\xi)}{b_1 \exp(\xi) + b_0 + b_{-1} \exp(-\xi)}. \quad (3.14)$$

Substituting Eq.(3.14) into Eq.(3.5), and by the help of Maple, we have

$$\frac{1}{A} [R_4 \exp(4\xi) + R_3 \exp(3\xi) + R_2 \exp(2\xi) + R_1 \exp(\xi) + R_0 + R_{-1} \exp(-\xi) + R_{-2} \exp(-2\xi) + R_{-3} \exp(-3\xi) + R_{-4} \exp(-4\xi)] = 0, \quad (3.15)$$



where

$$\begin{aligned}
 A &= (b_{-1} \exp(-\xi) + b_0 + b_1 \exp(\xi))^4, \\
 R_4 &= \xi_0 b_1^4, \\
 R_3 &= kca_1 b_1^2 b_0 - \frac{3}{4} n^2 a_0 b_1^3 + 4\xi_0 b_1^3 b_0 - \frac{1}{4} k^4 a_0 b_1^3 \\
 &\quad - kca_0 b_1^3 + \frac{3}{4} n^2 a_1 b_1^2 b_0 + \frac{1}{4} k^4 a_1 b_1^2 b_0, \\
 R_2 &= 2kca_1 b_1 b_0^2 + 2kca_1 b_1^2 b_{-1} - 2kca_0 b_0 b_1^2 - \frac{3}{2} k^3 a_1 b_0 a_0 b_1 \\
 &\quad - \frac{3}{2} n^2 a_{-1} b_1^3 + 6\xi_0 b_1^2 b_0^2 - 2k^4 a_{-1} b_1^3 + \frac{3}{4} k^3 a_1^2 b_0^2 + \frac{3}{4} k^3 a_0^2 b_1^2 \\
 &\quad + 4\xi_0 b_1^3 b_{-1} + 2k^4 a_1 b_1^2 b_{-1} + \frac{3}{2} n^2 a_1 b_1 b_0^2 + \frac{3}{2} n^2 a_1 b_1^2 b_{-1} \\
 &\quad - 2kca_{-1} b_1^3 - \frac{3}{2} n^2 a_0 b_1^2 b_0 - k^4 a_1 b_1 b_0^2 + k^4 a_0 b_1^2 b_0, \\
 R_1 &= 6kca_1 b_1 b_0 b_{-1} - \frac{1}{4} k^4 a_0 b_1 b_0^2 + 3k^3 a_1^2 b_0 b_{-1} - \frac{5}{4} k^4 a_{-1} b_1^2 b_0 \\
 &\quad + 3k^3 a_{-1} b_1^2 a_0 - \frac{3}{4} n^2 a_0 b_1 b_0^2 + kca_1 b_0^3 - \frac{3}{4} n^2 a_0 b_1^2 b_{-1} + 12\xi_0 b_1^2 b_0 b_{-1} \\
 &\quad + \frac{23}{4} k^4 a_0 b_1^2 b_{-1} - \frac{15}{4} n^2 a_{-1} b_1^2 b_0 - 3k^3 a_1 a_0 b_{-1} b_1 - kca_0 b_1 b_0^2 \\
 &\quad - 3k^3 a_1 b_0 a_{-1} b_1 - kca_0 b_1^2 b_{-1} - \frac{9}{2} k^4 a_1 b_1 b_0 b_{-1} + \frac{9}{2} n^2 a_1 b_1 b_0 b_{-1} \\
 &\quad - 5kca_{-1} b_1^2 b_0 + \frac{3}{4} n^2 a_1 b_0^3 + \frac{1}{4} k^4 a_1 b_0^3 + 4\xi_0 b_1 b_0^3, \\
 R_0 &= 4kca_1 b_1 b_{-1}^2 + 4kca_1 b_0^2 b_{-1} - 4kca_{-1} b_1^2 b_{-1} - 4kca_{-1} b_1 b_0^2 \\
 &\quad - 8k^4 a_1 b_1 b_{-1}^2 + k^4 a_1 b_0^2 b_{-1} + 8k^4 a_{-1} b_1^2 b_{-1} - k^4 a_{-1} b_1 b_0^2 \\
 &\quad - \frac{3}{2} k^3 a_1 b_0^2 a_{-1} + 6\xi_0 b_1^2 b_{-1}^2 + 3n^2 a_1 b_1 b_{-1}^2 + 3n^2 a_1 b_0^2 b_{-1} - 3n^2 a_{-1} b_1^2 b_{-1}, \\
 R_{-1} &= -\frac{9}{2} n^2 a_{-1} b_1 b_0 b_{-1} - 3k^3 a_{-1} b_1 a_0 b_{-1} + kca_0 b_1 b_{-1}^2 \\
 &\quad + \frac{9}{2} k^4 a_{-1} b_1 b_0 b_{-1} + kca_0 b_{-1} b_0^2 + 5kca_1 b_0 b_{-1}^2 - 3k^3 a_1 b_{-1} a_{-1} b_0 \\
 &\quad - \frac{3}{2} n^2 a_{-1} b_0^3 + 4\xi_0 b_0^3 b_{-1} - \frac{1}{4} k^4 a_{-1} b_0^3 + \frac{1}{4} k^4 a_0 b_{-1} b_0^2 + \frac{3}{4} n^2 a_0 b_{-1} b_0^2 \\
 &\quad + \frac{15}{4} n^2 a_1 b_0 b_{-1}^2 - kca_{-1} b_0^3 + 3k^3 a_1 b_{-1}^2 a_0 + 3k^3 a_{-1}^2 b_1 b_0 + 12\xi_0 b_1 b_0 b_{-1}^2 \\
 &\quad + \frac{3}{4} n^2 a_0 b_1 b_{-1}^2 + \frac{5}{4} k^4 a_1 b_0 b_{-1}^2 - \frac{23}{4} k^4 a_0 b_{-1}^2 b_1 - 6kca_{-1} b_1 b_0 b_{-1}, \\
 R_{-2} &= -\frac{3}{2} k^3 a_{-1} b_0 a_0 b_{-1} - 2kca_{-1} b_0^2 b_{-1} + 2kca_0 b_{-1}^2 b_0 - 2kca_{-1} b_1 b_{-1}^2 + 2k^4 a_1 b_{-1}^3 \\
 &\quad + \frac{3}{4} k^3 a_{-1}^2 b_0^2 + \frac{3}{4} k^3 a_0^2 b_{-1}^2 + 4\xi_0 b_1 b_{-1}^3 + \frac{3}{2} n^2 a_1 b_{-1}^3 + 6\xi_0 b_0^2 b_{-1}^2 - \frac{3}{2} n^2 a_{-1} b_0^2 b_{-1}, \\
 &\quad + \frac{3}{2} n^2 a_0 b_{-1}^2 b_0 - 2k^4 a_{-1} b_1 b_{-1}^2 + k^4 a_{-1} b_0^2 b_{-1} - k^4 a_0 b_{-1}^2 b_0 - \frac{3}{2} n^2 a_{-1} b_1 b_{-1}^2 + 2kca_1 b_{-1}^3, \\
 R_{-3} &= -kca_{-1} b_0 b_{-1}^2 + \frac{1}{4} k^4 a_0 b_{-1}^3 + 4\xi_0 b_0 b_{-1}^3 + \frac{3}{4} n^2 a_0 b_{-1}^3 \\
 &\quad - \frac{1}{4} k^4 a_{-1} b_0 b_{-1}^2 - \frac{3}{4} n^2 a_{-1} b_0 b_{-1}^2 + kca_0 b_{-1}^3, \\
 R_{-4} &= \xi_0 b_{-1}^4.
 \end{aligned}$$

(3.16)

Solving this system of algebraic equations by using Maple, we get the following results

Case 1: Consider

$$\begin{aligned}
 a_1 &= a_1, & a_0 &= \frac{b_0(a_1 - 2kb_1)}{b_1}, & a_{-1} &= 0, \\
 b_1 &= b_1, & b_0 &= b_0, & b_{-1} &= 0, \\
 c &= -\frac{3n^2 + k^4}{4k}, & k &= k, & \xi_0 &= 0,
 \end{aligned}$$



where  $b_0$  and  $b_1 \neq 0$  are free parameters. Substituting these results into (3.14), we get the following exact solution

$$u_1(x, y, t) = \frac{a_1 \exp\left(\frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+k^4)t^\alpha}{4k\Gamma(1+\alpha)}\right) + \frac{b_0(a_1-2kb_1)}{b_1}}{b_1 \exp\left(\frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+k^4)t^\alpha}{4k\Gamma(1+\alpha)}\right) + b_0}. \quad (3.18)$$

Case 2: Consider

$$\begin{aligned} a_1 &= 0, & a_0 &= \frac{b_0(a_{-1}+2kb_{-1})}{b_{-1}}, & a_{-1} &= a_{-1}, \\ b_1 &= 0, & b_0 &= b_0, & b_{-1} &= b_{-1}, \\ c &= -\frac{3n^2+k^4}{4k}, & k &= k, & \xi_0 &= 0, \end{aligned} \quad (3.19)$$

where  $b_{-1} \neq 0$  and  $a_{-1}$  are free parameters. Substituting these results into (3.14), we get the following exact solution

$$u_2(x, y, t) = \frac{\frac{b_0(a_{-1}+2kb_{-1})}{b_{-1}} + a_{-1} \exp\left(-\left(\frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+k^4)t^\alpha}{4k\Gamma(1+\alpha)}\right)\right)}{b_0 + b_{-1} \exp\left(-\left(\frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+k^4)t^\alpha}{4k\Gamma(1+\alpha)}\right)\right)}. \quad (3.20)$$

If we take  $b_1 = 1$ ,  $b_0 = 1$ ,  $a_1 = 1$  and  $k = 1$  Eq. (3.18)  $u_1$  becomes

$$u_1(x, y, t) = \frac{\cosh\left(\frac{x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) + \sinh\left(\frac{x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) - 1}{\cosh\left(\frac{x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) + \sinh\left(\frac{x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) + 1}, \quad (3.21)$$

Similarly,  $b_{-1} = 1$ ,  $b_0 = 1$ ,  $a_{-1} = 1$  and  $k = -1$  Eq. (3.20)  $u_2$  becomes

$$u_2(x, y, t) = \frac{\cosh\left(\frac{-x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} + \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) - \sinh\left(\frac{-x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} + \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) - 1}{\cosh\left(\frac{-x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} + \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) - \sinh\left(\frac{-x^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} + \frac{(3n^2+1)t^\alpha}{4\Gamma(1+\alpha)}\right) + 1}, \quad (3.22)$$

which are the exact solutions of the space-time fractional pKP equation.

**Remark 1:** Comparing our results, Eqs. (3.21) and (3.22), with Borhanifar's results in [38], it can be seen that the results are different. And these solutions have not been reported other authors in the literature.

Now we study the pKP equation to solve by using the  $\left(\frac{G'}{G}\right)$ -expansion method. By using the ansatz (3.5), for the linear term of highest order  $U'''$  with the highest order and the nonlinear term  $(U')^2$ , balancing  $U'''$  with  $(U')^2$  in Eq. (3.5) gives

$$m + 3 = 2m + 2, \quad (3.23)$$

so that

$$m = 1. \quad (3.24)$$



Suppose that the solutions of (3.5) can be expressed by a polynomial in  $\left(\frac{G'}{G}\right)$  as follows:

$$U(\xi) = a_0 + a_1 \left(\frac{G'}{G}\right), a_1 \neq 0. \tag{3.25}$$

By using Eq. (2.12) and Eq. (3.25) we have

$$U'(\xi) = -a_1 \left(\frac{G'}{G}\right)^2 - a_1 \lambda \left(\frac{G'}{G}\right) - a_1 \mu, \tag{3.26}$$

and

$$U'''(\xi) = -6a_1 \left(\frac{G'}{G}\right)^4 - 12a_1 \lambda \left(\frac{G'}{G}\right)^3 - (8a_1 \mu + 7a_1 \lambda^2) \left(\frac{G'}{G}\right)^2 - (8a_1 \lambda \mu + a_1 \lambda^3) \left(\frac{G'}{G}\right) - a_1 \mu^2 - 2a_1 \lambda^2 \mu. \tag{3.27}$$

By substituting Eqs. (3.25)-(3.27) into Eq. (3.5), collecting the coefficients of  $\left(\frac{G'}{G}\right)^i$  ( $i = 0, \dots, 4$ ) and setting them to zero, we obtain the equation system

$$\begin{aligned} -\frac{3}{2}k^4 a_1 + \frac{3}{4}k^3 a_1^2 &= 0, \\ -3k^4 a_1 \lambda + \frac{3}{2}k^3 a_1^2 \lambda &= 0, \\ -2k^4 a_1 \mu - \frac{7}{4}k^4 a_1 \lambda^2 + \frac{3}{2}k^3 a_1^2 \mu + \frac{3}{4}k^3 a_1^2 \lambda^2 - kca_1 - \frac{3}{4}n^2 a_1 &= 0, \\ -\frac{1}{4}k^4 a_1 \lambda^3 - \frac{3}{4}n^2 a_1 \lambda + \frac{3}{2}k^3 a_1^2 \lambda \mu - 2k^4 a_1 \lambda \mu - kca_1 \lambda &= 0, \\ -\frac{1}{2}k^4 a_1 \lambda^2 \mu - \frac{3}{4}n^2 a_1 \mu + \frac{3}{4}k^3 a_1^2 \mu^2 - \frac{1}{4}k^4 a_1 \mu^2 - kca_1 \mu + \xi_0 &= 0. \end{aligned} \tag{3.28}$$

By solving this system with the aid of Maple, we obtain

$$\begin{aligned} a_0 = a_0, \quad a_1 = 2k, \quad c = \frac{4k^4 \mu - k^4 \lambda^2 - 3n^2}{4k} \\ k = k, \quad n = n, \quad \xi_0 = \frac{1}{2}k^5 \lambda^2 \mu - \frac{1}{2}k^5 \mu^2, \end{aligned} \tag{3.29}$$

where  $\lambda$  and  $\mu$  are arbitrary constants. By using Eq. (3.29), expression (3.35) can be written as

$$U(\xi) = a_0 + 2k \left(\frac{G'}{G}\right). \tag{3.30}$$

By substituting general solutions of Eq. (2.12) into Eq. (3.30) we have three types of travelling wave solutions of the space-time fractional potential Kadomtsev–Petviashvili (pKP) equation as follows:

When  $\lambda^2 - 4\mu > 0$ , we obtain the hyperbolic function traveling wave solution

$$U_1(\xi) = a_0 - k\lambda + k\sqrt{\lambda^2 - 4\mu} \left( \frac{C_1 \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi + C_2 \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi}{C_1 \cosh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi + C_2 \sinh \frac{1}{2}\sqrt{\lambda^2 - 4\mu}\xi} \right), \tag{3.31}$$





where  $\xi = \frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} + \frac{(4k^4\mu - k^4\lambda^2 - 3n^2)t^\alpha}{4k\Gamma(1+\alpha)}$ , and  $C_1, C_2$  are arbitrary constants. In particular, if  $C_1 \neq 0, C_2 = 0, \lambda > 0, \mu = 0$ , then the traveling wave solution of (3.31) can be written as:

$$u_3(x, y, t) = a_0 - k\lambda + k\lambda \tanh \left\{ \frac{\lambda}{2} \left( \frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(k^4\lambda^2 + 3n^2)t^\alpha}{4k\Gamma(1+\alpha)} \right) \right\}. \quad (3.32)$$

And assuming  $C_1 = 0, C_2 \neq 0, \lambda > 0, \mu = 0$ , then we obtain

$$u_4(x, y, t) = a_0 - k\lambda + k\lambda \coth \left\{ \frac{\lambda}{2} \left( \frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(k^4\lambda^2 + 3n^2)t^\alpha}{4k\Gamma(1+\alpha)} \right) \right\}. \quad (3.33)$$

When  $\lambda^2 - 4\mu < 0$ , we obtain the trigonometric function traveling wave solution

$$U_2(\xi) = a_0 - k\lambda + ik\lambda \left( \frac{-C_1 \sin \frac{1}{2} \sqrt{4\mu - \lambda^2} \xi + C_2 \cos \frac{1}{2} \sqrt{4\mu - \lambda^2} \xi}{C_1 \cos \frac{1}{2} \sqrt{4\mu - \lambda^2} \xi + C_2 \sin \frac{1}{2} \sqrt{4\mu - \lambda^2} \xi} \right), \quad (3.34)$$

Also, if we assume  $C_1 \neq 0, C_2 = 0, \lambda > 0, \mu = 0$ , then

$$u_5(x, y, t) = a_0 - k\lambda + k\lambda \tanh \left\{ \frac{\lambda}{2} \left( \frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(k^4\lambda^2 + 3n^2)t^\alpha}{4k\Gamma(1+\alpha)} \right) \right\}, \quad (3.35)$$

and when  $C_1 = 0, C_2 \neq 0, \lambda > 0, \mu = 0$ , the solution of Eq. (3.34) becomes

$$u_6(x, y, t) = a_0 - k\lambda + k\lambda \coth \left\{ \frac{\lambda}{2} \left( \frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} - \frac{(k^4\lambda^2 + 3n^2)t^\alpha}{4k\Gamma(1+\alpha)} \right) \right\}. \quad (3.36)$$

So we obtain the solutions  $u_3(x, y, t)$  and  $u_4(x, y, t)$ .

When  $\lambda^2 - 4\mu = 0$ , we obtain the rational function solution

$$u_7(x, y, t) = a_0 - k\lambda + \frac{2kC_2}{C_1 + C_2 \left( \frac{kx^\alpha}{\Gamma(1+\alpha)} + \frac{ny^\alpha}{\Gamma(1+\alpha)} + \frac{(4k^4\mu - k^4\lambda^2 - 3n^2)t^\alpha}{4k\Gamma(1+\alpha)} \right)}. \quad (3.37)$$

**Remark 2:** Compare our results, Eqs. (3.32), (3.33) and (3.37), with Budhiraja's and Zayed's solutions in [39, 40], it can be seen that the results are different.



#### 4. CONCLUSIONS

In this paper, we have seen that three types of exact analytical solutions including the hyperbolic function solutions, trigonometric function solutions and rational solutions for the space-time fractional pKP equation are successfully found out by using the exp-function and  $(G'/G)$ -expansion methods. This study shows that the exp-function and  $(G'/G)$ -expansion methods are quite efficient and practically well suited for finding exact solutions of the pKP equation. The performance of these methods are reliable and effective and give the exact solitary wave solutions and periodic wave solutions. The availability of computer symbolic systems like Mathematica or Maple facilitates the tedious algebraic calculations. Thus, we deduce that the proposed method can be extended to solve many systems of nonlinear time-fractional partial differential equations.

#### REFERENCES

- [1] Miller, K. S., Ross, B., An Introduction to the Fractional Calculus and Fractional Differential Equations, Wiley, New York (1993)
- [2] Podlubny, I., Fractional Differential Equations, Academic Press, California (1999)
- [3] Kilbas, A. A., Srivastava, H. M., Trujillo, J. J., Theory and Applications of Fractional Differential Equations, Elsevier, Amsterdam (2006)
- [4] Zheng, B.,  $(G'/G)$ -expansion method for solving fractional partial differential equations in the theory of mathematical physics, Commun. Theor. Phys., 58 (2012) 623–630
- [5] Gepreel, K. A., Omran, S., Exact solutions for nonlinear partial fractional differential equations, Chin. Phys. B, 21 (2012) 110204
- [6] Bekir, A., Güner, Ö., Exact solutions of nonlinear fractional differential equations by  $(G'/G)$ -expansion method, Chin. Phys. B, 22, 11 (2013) 110204
- [7] Shang, N., Zheng, B., Exact Solutions for Three Fractional Partial Differential Equations by the  $(G'/G)$  Method, International Journal of Applied Mathematics, 43 (2013) 3
- [8] Zhang, S., Zong Q-A., Liu, D., Gao, Q., A Generalized Exp-Function Method for Fractional Riccati Differential Equations, Communications in Fractional Calculus, 1 (2010) 48-51.
- [9] Bekir, A., Güner, Ö., Cevikel, A.C., Fractional Complex Transform and exp-Function Methods for Fractional Differential Equations, Abstract and Applied Analysis, 2013, (2013) 426462
- [10] Zheng, B., Exp-Function Method for Solving Fractional Partial Differential Equations, The Scientific World Journal, 2013 (2013) 465723
- [11] Bekir, A., Güner, Ö., Cevikel, A.C., Using A Complex Transformation with Exp-Function Method to get an Exact Solutions for Fractional Differential Equation, Current Advances in Mathematics, 2,1 (2014) 22-31
- [12] Lu, B., The first integral method for some time fractional differential equations, J. Math. Anal. Appl., 395 (2012) 684–693
- [13] Eslami, M., Vajargah, B. F., Mirzazadeh, M., Biswas, A., Application of first integral method to fractional partial differential equations, Indian J. Phys. 88, 2 (2014) 177–184
- [14] Zhang, S., Zhang, H-Q., Fractional sub-equation method and its applications to nonlinear fractional PDEs, Physics Letters A, 375 (2011) 1069–1073
- [15] Zheng, B., Wen, C., Exact solutions for fractional partial differential equations by a new fractional sub-equation method, Advances in Difference Equations, 2013 (2013) 199
- [16] Alzaidy, J. F., Fractional Sub-Equation Method and its Applications to the Space–Time Fractional Differential Equations in Mathematical Physics, British Journal of Mathematics & Computer Science, 3 (2013) 153-163
- [17] Liu, W., Chen, K., The functional variable method for finding exact solutions of some nonlinear time-fractional differential equations, Pramana–J. Phys., 81, 3 (2013) 377–384



- [18] Bulut, H., Baskonus, H M., Pandir, Y., The Modified Trial Equation Method for Fractional Wave Equation and Time Fractional Generalized Burgers Equation, *Abstract and Applied Analysis*, 2013 (2013) 636802
- [19] Pandir, Y., Gurefe, Y., Misirli, E., New Exact Solutions of the Time-Fractional Nonlinear Dispersive KdV Equation, *International Journal of Modeling and Optimization*, 3 (2013) 4
- [20] Pandir, Y., Gurefe, New exact solutions of the generalized fractional Zakharov-Kuznetsov equations, *Life Science Journal*, 10, 2 (2013) 2701–2705
- [21] Taghizadeh N, Mirzazadeh M, Rahimian M, Akbari M., Application of the simplest equation method to some time-fractional partial differential equations, *Ain Shams Engineering Journal*, 4 (2013) 897–902
- [22] Jumarie, G., Modified Riemann–Liouville derivative and fractional Taylor series of nondifferentiable functions further results, *Comput. Math. Appl.*, 51 (2006) 1367–1376
- [23] Jumarie, G., Table of some basic fractional calculus formulae derived from a modified Riemann–Liouville derivative for nondifferentiable functions, *Appl. Maths. Lett.*, 22 (2009) 378–385
- [24] Li, Z.B., He, J H. Fractional complex transform for fractional differential equations, *Math. Comput. Appl.*, 15 (2010) 970–973
- [25] Li, Z.B., He, J. H., Application of the fractional complex transform to fractional differential equations *Nonlinear Sci. Lett. A Math. Phys. Mech.*, 2 (2011)121-126
- [26] He, J.H., Abdou, M.A., New periodic solutions for nonlinear evolution equations using Exp-function method, *Chaos Solitons Fractals*, 34 5 (2007) 1421-1429
- [27] He, J.H., Wu, X.H., Exp-function method for nonlinear wave equations, *Chaos Solitons Fractals*, 30 (2006) 700-708
- [28] Bekir, A., Boz, A. Application of He’s exp-function method for nonlinear evolution equations, *Computers and Mathematics with Applications*, 58 (2009) 2286-2293
- [29] Wang, M., Li, X., Zhang, J., The  $(G'/G)$ -expansion method and traveling wave solutions of nonlinear evolution equations in mathematical physics, *Phys. Lett. A*, 372 (2008) 417- 423
- [30] Bekir, A., Application of the  $(G'/G)$ -expansion method for nonlinear evolution equations, *Physics Letters, A* 372 (2008) 3400-3406
- [31] Alzaidy, J. F., The Fractional Sub-Equation Method and Exact Analytical Solutions for Some Nonlinear Fractional PDEs, *American Journal of Mathematical Analysis*, 1 1 (2013) 14-19
- [32] Biswas, A., Ranasinghe, A., 1-Soliton solution of Kadomtsev-Petviashvili equation with power law nonlinearity, *Applied Mathematics and Computation*, 214, 2 (2009) 645-647
- [33] Biswas, A., Ranasinghe, A., Topological 1-Soliton solution of Kadomtsev-Petviashvili equation with power law nonlinearity, *Applied Mathematics and Computation*, 217, 4 (2010) 1771-1773
- [34] Jawad, A.J.M., Petkovic, M., Biswas, A., Soliton solutions for nonlinear Calogero-Degasperis and potential Kadomtsev-Petviashvili equations, *Computers and Mathematics with Applications*, 62, 6 (2011) 2621-2628
- [35] Triki, H., Sturdevant, B.J.M., Omar, T.H, Aldossary, M., Biswas, A., Shock wave solutions of the variants of Kadomtsev-Petviashvili equation, *Canadian Journal of Physics*, 89, 9 (2011) 979-984
- [36] Bhrawy, A.H., Abdelkawy, M.A., Kumar, S., Biswas, A., Solitons and other solutions to Kadomtsev-Petviashvili equation of B-type, *Romanian Journal of Physics*, 58, 7-8 (2013) 729-748
- [37] Ebadi, G., Fard, N.Y., Bhrawy, A.H., Kumar, S., Triki, H., Yildirim, A., Biswas, A., Solitons and other solutions to the (3+1)-dimensional extended Kadomtsev-Petviashvili equation with power law nonlinearity, *Romanian Reports in Physics*, 65, 1 (2013) 27-62
- [38] Borhanifar, A., Kabir, M.M., New periodic and soliton solutions by application of Exp-function method for nonlinear evolution equations, *J. Comput. Appl. Math.*, 229 (2009) 158-167
- [39] Budhiraja, R., Gupta, R.K., Bhatia, S.S., The New Generalized  $(G'/G)$ -expansion Method for Solving (2+1) Dimensional PKP Equation, *International Journal of Nonlinear Science*, 14 1 (2012) 48-52
- [40] Zayed, E.M.E., Amer, Y.A., Shohib, R.M.A., Exact traveling wave solutions for nonlinear fractional partial differential equations using the improved  $(G'/G)$ -expansion method, *International Journal of Engineering and Applied Science*, 4, 7 (2014) 18-31

