



Existence of positive solution to a class of boundary value problems of fractional differential equations

Amjad Ali*

Department of Mathematics, University of Malakand,
Chakadara Dir(L), Khyber Pakhtunkhwa, Pakistan.
E-mail: amjadalimna@yahoo.com

Kamal Shah

Department of Mathematics, University of Malakand,
Chakadara Dir(L), Khyber Pakhtunkhwa, Pakistan.
E-mail: kamalshah408@gmail.com

Rahmat Ali Khan

Department of Mathematics, University of Malakand,
Chakadara Dir(L), Khyber Pakhtunkhwa, Pakistan.
E-mail: rahmat.alipk@yahoo.com

Abstract

This paper is devoted to the study of establishing sufficient conditions for existence and uniqueness of positive solution to a class of non-linear problems of fractional differential equations. The boundary conditions involved Riemann-Liouville fractional order derivative and integral. Further, the non-linear function f contain fractional order derivative which produce extra complexity. Thank to classical fixed point theorems of nonlinear alternative of Leray-Schauder and Banach Contraction principle, sufficient conditions are developed under which the proposed problem has at least one solution. An example has been provided to illustrate the main results.

Keywords. Boundary value problem, Existence and uniqueness results, Fractional differential differential equations, Classical fixed point theorem..

2010 Mathematics Subject Classification. 26A33, 34A37, 34B05.

1. INTRODUCTION

Fractional differential equations have many applications in various field of science and technology. The important applications, we may observe in various scientific and engineering disciplines like, physics, chemistry, biology, photoelasticity, control theory and signal processing. Moreover, many applications are found in dynamics, aerodynamics, electrostatics and biophysics, economics, polymers rheology, thermodynamics and biomedical science, (see [9, 11, 14, 15, 19, 26, 25]). Due to the aforementioned applications, the concerned subject has got much attention from the researchers of mathematics, physics etc. The researchers has been studied by details its theory of existences, numerical analysis and qualitative analysis. Plenty of work can be found on the said area, for more detail, we refer the reader to [1, 2, 3, 13, 7, 6, 17, 24, 26]. It

Received: 15 July 2016 ; Accepted: 7 November 2016.

* Corresponding author.

is to be noted that Caputo fractional derivatives plays a vital role in the modeling of dynamical and physical problems as it has known physical interpretation for initial and boundary value problems like classical differential equations. But on the other hand derivative in Riemann-Liouville sense also plays important role in the area of applied analysis. Existence theory for real world problems carried out by researchers of mathematics for fractional differential equations with boundary conditions has been explored and many research articles can be found regarding this topic, for detail (see [8, 4, 18, 28, 10, 20, 27, 21, 22, 23]). In [5], Bai studied existence and multiplicity of positive solutions to the following boundary value problem

$$\begin{aligned} \mathcal{D}^\alpha u(t) + a(t)f(t, u'(t)) &= 0; \quad 0 < t < 1, n-1 \leq \alpha < n, n > 2, \\ u(t)|_{t=0} = u'(t)|_{t=0} = u''(t)|_{t=0} = u''(t)|_{t=1} &= 0, \end{aligned}$$

where $f : [0, 1] \times [0, \infty) \rightarrow (-\infty, +\infty)$ is nonlinear continuous function and \mathcal{D}^α is the Riemann-Liouville fractional derivative. In [16], Kosmatov studied existence and uniqueness of solutions to two-point boundary value problem for fractional differential equations of the form

$$\begin{aligned} \mathcal{D}^\alpha u(t) + f(t, u(t), u'(t)) &= 0; \quad t \in (0, 1), 1 < \alpha < 2, \\ u(t)|_{t=0} = u(t)|_{t=1} &= 0, \end{aligned}$$

where \mathcal{D}^α is the Riemann-Liouville fractional derivative. In these cited papers, the nonlinearity f involved the classical order derivative. The case where the nonlinearity f explicitly depends of fractional order derivative is important theoretically as well as in application point of view and requires more efforts to study existence results. Goodrich [12], proved multiple solutions by using Schauder's fixed point theorem for fractional differential equation of the form

$$\begin{aligned} -\mathcal{D}^\alpha u(t) &= f(t, u(t)); \quad 0 < t < 1, n-1 < \alpha \leq n, n \in \mathbb{N}, \\ u^{(i)}(t)|_{t=0} = \mathcal{D}^\beta u(t)|_{t=1} &= 0, \end{aligned}$$

where $0 \leq i \leq n-2$, $1 \leq \beta \leq n-2$, $\alpha > 3$. The nonlinear function $f : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ is continuous.

Motivated by the above mentioned work, this paper is concerned to establish conditions for existence and uniqueness of solution to fractional order differential equation with boundary condition provided by

$$\begin{aligned} \mathcal{D}^\alpha u(t) &= f(t, u(t), \mathcal{D}^p u(t)) = h(t); \quad 2 < \alpha < 3, 0 < p < 1, \\ \mathcal{I}^{3-\alpha} u(t)|_{t=0} = \mathcal{D}^{\alpha-2} u(t)|_{t=0} = u(t)|_{t=1} &= 0. \end{aligned} \tag{1.1}$$

The required conditions are obtained by using classical fixed point theorem like non-linear alternative of Leray-Schauder type and Banach contraction theorem. In last, the paper is enriched by providing a suitable example to illustrate the main results. The paper is organized: In second section, some preliminary results are provided for obtaining the main results. In third Section, we provided main results in which derivation of the Green's function, integral formulations of the problem, necessary and sufficient conditions for the existence and uniqueness of solutions by applying



some classical fixed point theorems. Section 4 is devoted to illustrate the main results by an example.

2. PRELIMINARIES

We recall some basic definitions and known results from fractional calculus, functional analysis, which can be found in [14, 26, 21, 28, 27].

Definition 2.1. The fractional integral of order $q \in \mathbb{R}_+$ of the function $u : (0, \infty) \rightarrow R$ is defined as

$$\mathcal{I}^\alpha u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} u(s) ds,$$

where the integral on the right side is pointwise defined on $(0, \infty)$.

Definition 2.2. The Riemann-Liouville fractional order derivative of a function u on the interval $[a, b]$ is defined by

$$\mathcal{D}^\alpha u(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt}\right)^n \int_a^t (t-s)^{n-\alpha-1} u(s) ds,$$

where $n = [\alpha] + 1$. It is to be noted that integral on the right hand side is pointwise defined on $(0, \infty)$.

The following Lemmas are necessary in this paper.

Lemma 2.3. [17] *The fractional order differential equation of order $\alpha > 0$ of the form*

$$\mathcal{D}^\alpha u(t) = 0, \quad n-1 < \alpha \leq n,$$

has a unique solution of the form

$$u(t) = C_1 t^{\alpha-1} + C_2 t^{\alpha-2} + \dots + C_n t^{\alpha-n}, \quad \text{where } C_i \in R,$$

$i = 1, 2, \dots, n$, and $n = [\alpha] + 1$, where $[\alpha]$ represents integer part of α .

Lemma 2.4. [17] *The following result holds for a fractional derivative and integral of order α*

$$I^\alpha \mathcal{D}^\alpha u(t) = u(t) + C_1 t^{\alpha-1} + C_2 t^{\alpha-2} + \dots + C_n t^{\alpha-n},$$

where $C_i \in R$, $i = 1, 2, \dots, n$, and $n = [\alpha] + 1$, where $[\alpha]$ represents integer part of α .

To establish, the main results, we need the following Theorems in this paper.

Theorem 2.5. *(The nonlinear alternative of Leray-Schauder type [27]) Let D be a closed convex subset of a Banach space X . Consider a relative open subset C of D such that $0 \in C$ and let $T : C \rightarrow D$ be continuous and compact mapping. Then either*

- (1) *the mapping T has a fixed point in C ; or*
- (2) *there exist $u \in \partial C$ and $\lambda \in (0, 1)$ with $u = \lambda T u$.*



Theorem 2.6. (*The Banach contraction theorem* [27]) *Let D be a closed subset of a Banach space X and $T : D \rightarrow D$. Then T has a unique fixed point u in D and for any initial value $u_0 \in D$, the successive approximation converges to u_0 if*

$$\|Tu - T\bar{u}\| \leq k\|u - \bar{u}\|, \text{ for all } u, \bar{u} \in D \text{ with } 0 < k < 1.$$

3. MAIN RESULTS

Theorem 3.1. *Let $h(t) \in C[0, 1]$, then the boundary value problem*

$$\begin{aligned} \mathcal{D}^\alpha u(t) &= h(t), 2 < \alpha < 3, t \in [0, 1], \\ \mathcal{I}^{3-\alpha} u(t)|_{t=0} &= \mathcal{D}^{\alpha-2} u(t)|_{t=0} = u(t)|_{t=1} = 0, \end{aligned} \quad (3.1)$$

has a unique solution given by

$$u(t) = \int_0^1 G(t, s)h(s)ds,$$

where $G(t, s)$ is the Green's function given as

$$G(t, s) = \frac{1}{\Gamma(\alpha)} \begin{cases} (t-s)^{\alpha-1} - t^{\alpha-1}(1-s)^{\alpha-1}, & 0 \leq s \leq t \leq 1; \\ -t^{\alpha-1}(1-s)^{\alpha-1}, & 0 \leq t \leq s \leq 1. \end{cases}$$

Proof. In view of Lemma(2.4), boundary value problem(3.1) is written as

$$u(t) = C_1 t^{\alpha-1} + C_2 t^{\alpha-2} + C_3 t^{\alpha-3} + \mathcal{I}^\alpha h(t). \quad (3.2)$$

Now for boundary conditions, (3.2) implies that

$$\begin{aligned} \mathcal{I}^{3-\alpha} u(t) &= C_1 \mathcal{I}^{3-\alpha} t^{\alpha-1} + C_2 \mathcal{I}^{3-\alpha} t^{\alpha-2} + C_3 \mathcal{I}^{3-\alpha} t^{\alpha-3} + \mathcal{I}^{3-\alpha} \mathcal{I}^\alpha h(t) \\ &= C_1 \frac{\Gamma(\alpha-1+1)}{\Gamma(3-\alpha+\alpha-1+1)} t^{3-\alpha+\alpha-1} + C_2 \frac{\Gamma(\alpha-2+1)}{\Gamma(3-\alpha+\alpha-2+1)} t^{3-\alpha+\alpha-2} \\ &\quad + C_3 \frac{\Gamma(\alpha-3+1)}{\Gamma(3-\alpha+\alpha-3+1)} t^{3-\alpha+\alpha-3} + \mathcal{I}^3 h(t) \\ \mathcal{I}^{3-\alpha} u(t) &= C_1 \frac{\Gamma(\alpha)}{\Gamma(3)} t^2 + C_2 \frac{\Gamma(\alpha-1)}{\Gamma(2)} t + C_3 \frac{\Gamma(\alpha-2)}{\Gamma(1)} + \mathcal{I}^3 h(t). \end{aligned} \quad (3.3)$$

In view of $\mathcal{I}^{3-\alpha} u(t)|_{t=0} = 0$ and also we have $\mathcal{I}^3 h(t) \rightarrow 0$ as $t \rightarrow 0$, in(3.3), we get $C_3 = 0$. Now for boundary conditions $\mathcal{D}^{\alpha-2} u(t)|_{t=0} = 0$, (3.2) implies that

$$\begin{aligned} \mathcal{D}^{\alpha-2} u(t) &= \mathcal{D}^{\alpha-2} \left[C_1 t^{\alpha-1} + C_2 t^{\alpha-2} + \mathcal{I}^\alpha h(t) \right] \\ &= C_1 \frac{\Gamma(\alpha)}{\Gamma(\alpha-1-\alpha+2+1)} t^{\alpha-1-\alpha+2} + C_2 \frac{\Gamma(\alpha-1)}{\Gamma(\alpha-2-\alpha+2+1)} t^{\alpha-2-\alpha+2} + \mathcal{I}^2 h(t) \\ &= C_1 \frac{\Gamma(\alpha)}{\Gamma(2)} t + C_2 \frac{\Gamma(\alpha-1)}{\Gamma(1)} + \mathcal{I}^2 h(t) \end{aligned} \quad (3.4)$$



which on application of $\mathcal{D}^{\alpha-2}u(t)|_{t=0} = 0$, and $\mathcal{I}^2h(t) \rightarrow 0$ as $t \rightarrow 0$, we get $C_2 = 0$. Hence (3.2) implies that

$$u(t) = C_1 t^{\alpha-1} + \mathcal{I}^\alpha h(t). \tag{3.5}$$

Using $u(t)|_{t=1} = 0$ in (3.5), we get $C_1 = -\mathcal{I}^\alpha h(1)$. Thus (3.2) becomes

$$\begin{aligned} u(t) &= -t^{\alpha-1}\mathcal{I}^\alpha h(1) + \mathcal{I}^\alpha h(t) \\ &= \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} h(s) ds - \frac{t^{\alpha-1}}{\Gamma(\alpha)} \int_0^1 (1-s)^{\alpha-1} h(s) ds \\ &= \int_0^1 G(t,s) h(s) ds. \end{aligned}$$

Hence the BVP (1.1) is equivalent to the following integral equation

$$u(t) = \int_0^1 G(t,s) f(s, u(s), \mathcal{D}^p u(s)) ds. \tag{3.6}$$

□

Lemma 3.2. *The Green's function $G(t, s)$ defined in Theorem 3.1 satisfies the following inequality*

$$\int_0^1 |G(t, s)| ds \leq \frac{2}{\Gamma(\alpha + 1)}. \tag{3.7}$$

Proof. Proof is easy, so we omit it. □

Assume that the following hold,

(A₁) For any $x, y, u, v \in R$, there exists $L > 0$ such that

$$|f(t, u, v) - f(t, x, y)| \leq L(|u - x| + |v - y|);$$

(A₂) $f : I \times R^2 \rightarrow R$ is continuous;

(A₃) There exists $a_0(t) \in L^1[0, 1]$ such that $|a_0(t)| \leq \delta$ and $b, c \geq 0$ with

$$|f(t, u, v)| \leq a_0(t) + b|u|^\rho + c|v|^\theta,$$

where $0 < \rho < \theta < 1$; or

$$|f(t, u, v)| \leq b|u|^\rho + c|v|^\theta \text{ for } \rho, \theta \geq 1, b, c > 0.$$

Throughout this paper, we use the following space defined by

$$\begin{aligned} X &= \{u(t) | u(t) \in C^1[0, 1] \text{ and } \mathcal{D}^p u(t) \in C^1[0, 1], 0 < p < 1\} \\ \|u\| &= \max_{t \in [0, 1]} |u(t)| + \max_{t \in [0, 1]} |\mathcal{D}^p u(t)|, \end{aligned}$$

then $(X, \|\cdot\|)$ is Banach space. Let define an operator $T : X \rightarrow X$ by

$$Tu(t) = \int_0^1 G(t,s) f(s, u(s), \mathcal{D}^p u(s)) ds, \quad t \in [0, 1]. \tag{3.8}$$



Theorem 3.3. *Under the assumptions (A_1) to (A_3) BVP (1.1) has at least one solution.*

Proof. Let us define $T : X \rightarrow X$ by

$$Tu(t) = \int_0^1 G(t, s) f(s, u(s), \mathcal{D}^p u(s)) ds.$$

Define a closed set as

$$B = \{u(t) | u \in X : \|u\| \leq R, 0 \leq t \leq 1\}, \text{ where } \max \left\{ (3Ab)^{\frac{1}{3}-\rho}, (3Ac)^{\frac{1}{3}-\theta}, 3k \right\} \leq R,$$

where

$$A = \frac{2\alpha - p}{\alpha\Gamma(\alpha - p + 1)} + \frac{2}{\Gamma(\alpha + 1)}$$

$$k = \delta \left[\frac{2}{\Gamma(\alpha + 1)} + \frac{1}{\alpha\Gamma(\alpha - p)} + \frac{1}{\Gamma(\alpha - p + 1)} \right].$$

For any $u \in B$ and $T : B \rightarrow B$, we have

$$\begin{aligned} |Tu(t)| &= \left| \int_0^1 G(t, s) f(s, u(s), \mathcal{D}^p u(s)) ds \right| \\ &\leq \int_0^1 |G(t, s)| |f(s, u(s), \mathcal{D}^p u(s))| ds \\ &\leq \int_0^1 |G(t, s)| (|a_0(s)| + b|u(s)|^\rho + c|\mathcal{D}^p u(s)|^\theta) ds \tag{3.9} \\ &\leq \int_0^1 |G(t, s)| |a_0(s)| ds + (bR^\rho + cR^\theta) \int_0^1 |G(t, s)| ds \\ &\leq \frac{2}{\Gamma(\alpha + 1)} \left[\delta + bR^\rho + cR^\theta \right]. \end{aligned}$$

Now from (3.6), we have

$$\begin{aligned} \mathcal{D}^p u(t) &= \mathcal{D}^p (-\mathcal{I}^\alpha h(1)t^{\alpha-1} + \mathcal{I}^\alpha h(t)) = -\mathcal{I}^\alpha h(1)\mathcal{D}^p t^{\alpha-1} + \mathcal{D}^p \mathcal{I}^\alpha h(t) \\ &= -\mathcal{I}^\alpha h(1) \frac{\Gamma(\alpha)t^{\alpha-1-p}}{\Gamma(\alpha-p)} + \mathcal{I}^{\alpha-\beta} h(t). \end{aligned}$$



Thus by applying operator T , we have

$$\begin{aligned}
 |\mathcal{D}^p Tu(t)| &= \left| -\frac{t^{\alpha-p-1}}{\Gamma(\alpha-p)} \int_0^1 (1-s)^{\alpha-1} f(s, u(s), \mathcal{D}^p u(s)) ds + \frac{1}{\Gamma(\alpha-p)} \int_0^t (t-s)^{\alpha-p-1} f(s, u(s), \mathcal{D}^p u(s)) ds \right| \\
 &\leq \frac{1}{\Gamma(\alpha-p)} \left\{ \left[\int_0^1 (1-s)^{\alpha-1} (|a_0(s)| + bR^\rho + cR^\theta) ds \right] \right. \\
 &\quad \left. + \left[\int_0^t (t-s)^{\alpha-p-1} (|a_0(s)| + bR^\rho + cR^\theta) ds \right] \right\} \\
 &\leq \frac{\delta}{\Gamma(\alpha-p)} \left[\frac{1}{\alpha} + \frac{1}{\alpha-p} \right] + \frac{2\alpha-p}{\alpha\Gamma(\alpha-p+1)} (bR^\rho + cR^\theta).
 \end{aligned}$$

Which implies that

$$|\mathcal{D}^p Tu(t)| \leq \frac{\delta}{\Gamma(\alpha-p)} \left[\frac{1}{\alpha} + \frac{1}{\alpha-p} \right] + \frac{2\alpha-p}{\alpha\Gamma(\alpha-p+1)} (bR^\rho + cR^\theta). \quad (3.10)$$

Adding (3.9) and (3.10), using $|a_0(s)| \leq \delta$, we obtain

$$\begin{aligned}
 |Tu(t)| + |\mathcal{D}^p Tu(t)| &\leq \frac{2\delta}{\Gamma(\alpha+1)} + \frac{2(bR^\rho + cR^\theta)}{\Gamma(\alpha+1)} + \frac{\delta}{\Gamma(\alpha-p)} \left[\frac{1}{\alpha} + \frac{1}{\alpha-p} \right] \\
 &\quad + \frac{2\alpha-p}{\alpha\Gamma(\alpha-p+1)} (bR^\rho + cR^\theta) \\
 &\leq \delta \left[\frac{2}{\Gamma(\alpha+1)} + \frac{1}{\alpha\Gamma(\alpha-p)} + \frac{1}{\Gamma(\alpha-p+1)} \right] \\
 &\quad + \left[\frac{2\alpha-p}{\alpha\Gamma(\alpha-p+1)} + \frac{2}{\Gamma(\alpha+1)} \right] (bR^\rho + cR^\theta)
 \end{aligned}$$

which implies that $\|Tu\| \leq k + (bR^\rho + cR^\theta)A \leq \frac{R}{3} + \frac{R}{3} + \frac{R}{3} = R$.

Also $Tu(t)$ and $\mathcal{D}^p Tu(t)$ are continuous. Thus $T : B \rightarrow B$ is well defined and bounded. Similarly using same argument for other condition of assumption (A_3) , we get

$$\|Tu\| < \frac{R}{2} + \frac{R}{2} = R.$$

Now, we are going to show that T is completely continuous operator. Let $\tau \leq t \in [0, 1]$, then

$$\begin{aligned}
 |Tu(t) - Tu(\tau)| &= \left| \int_0^1 (G(t, s) - G(\tau, s)) f(s, u, \mathcal{D}^p u) ds \right| \\
 &\leq \int_0^1 |G(t, s) - G(\tau, s)| |f(s, u, \mathcal{D}^p u)| ds. \quad (3.11)
 \end{aligned}$$



Let $c, d \in (\tau, t)$ for $0 \leq s \leq t$ and by the application of Lagrange's mean value theorem, we proceed as

$$\begin{aligned} |G(t, s) - G(\tau, s)| &= \frac{1}{\Gamma(\alpha)} [(t-s)^{\alpha-1} - (\tau-s)^{\alpha-1} + (1-s)^{\alpha-1}(\tau^{\alpha-1} - t^{\alpha-1})] \\ &\leq \frac{\alpha-1}{\Gamma(\alpha)} [(c-s)^{\alpha-2}(t-\tau) - (1-s)^{\alpha-1}d^{\alpha-2}(t-\tau)]. \end{aligned}$$

Similarly there exists $e \in (t, \tau)$ for $0 \leq t \leq s$

$$(\tau^{\alpha-1} - t^{\alpha-1}) \leq (\alpha-1)e^{\alpha-2}(\tau-t)(1-s)^{\alpha-1}.$$

Since $\alpha \leq 3$, therefore using $\alpha-1 \leq 2$, then (3.11) implies that

$$\begin{aligned} |Tu(\tau) - Tu(t)| &\leq \frac{(t-\tau)}{\Gamma(\alpha)} \int_0^1 2\{(c-s)^{\alpha-1} - d^{\alpha-2}(1-s)^{\alpha-1} - e^{\alpha-2}(1-s)^{\alpha-1}\}(a_0(s) + bR^\rho + cR^\theta) ds. \end{aligned} \quad (3.12)$$

Let

$$\mathbf{G}_{c,d,e}(s) = 2\{(c-s)^{\alpha-2} - (1-s)^{\alpha-1}d^{\alpha-2} - e^{\alpha-2}(1-s)^{\alpha-1}\}.$$

Then (3.12) becomes

$$|Tu(t) - Tu(\tau)| \leq \frac{(t-\tau)}{\Gamma(\alpha)} \int_0^1 \mathbf{G}_{c,d,e}(s)(a_0(s) + bR^\rho + cR^\theta) ds. \quad (3.13)$$

Similarly one has

$$|D^p Tu(t) - D^p Tu(\tau)| \leq \frac{(t-\tau)}{\Gamma(\alpha)} \int_0^1 \mathbf{G}_{c,d,e}(s)(a_0(s) + bR^\rho + cR^\theta) ds \quad (3.14)$$

Now from (3.13) and (3.14), we have

$$\|Tu(t) - Tu(\tau)\| \leq \frac{(t-\tau)}{\Gamma(\alpha)} \left[\int_0^1 \mathbf{G}_{c,d,e}(s) + \mathbf{G}_{c,d,e}(s) \right] [(a_0(s) + bR^\rho + cR^\theta) ds]. \quad (3.15)$$

Now $t \rightarrow \tau$ in (3.15) gives

$$\|Tu(t) - Tu(\tau)\| \rightarrow 0.$$

Thus T is equicontinuous, as T is uniformly continuous on $[0, 1]$. Thus $T(B)$ is equicontinuous set and also is bounded. Moreover, $T(B) \subseteq B$. It follows that T is completely continuous.

Thus T is at least one fixed point by using nonlinear alternative of Leray-Schauder theorem 2.5, which is the corresponding solution of (1.1). This completes the proof. \square

Theorem 3.4. *If $(A_1), (A_2)$ hold and $L\Omega = \left(\frac{2}{\Gamma(\alpha+1)} + \frac{1}{\Gamma(\alpha-p+1)}\right) L < 1$. Then T has a unique fixed point.*



Proof. Let $u(t), v(t) \in C^1[0, 1]$, then we have

$$|Tu(t) - Tv(t)| \leq \int_0^1 |G(t, s)| |f(t, u, \mathcal{D}^p u) - f(t, v, \mathcal{D}^p v)| ds$$

$$\leq \max \int_0^1 \left(\frac{(t-s)^{\alpha-1} - t^{\alpha-1}(1-s)^{\alpha-1}}{\Gamma(\alpha)} \right) L(|u-v| + |\mathcal{D}^p u - \mathcal{D}^p v|) ds.$$

Thus $\|Tu - Tv\| \leq \frac{2L}{\Gamma(\alpha + 1)} \|u - v\|.$ (3.16)

Similarly

$$\|\mathcal{D}^p Tu - \mathcal{D}^p Tv\| \leq \frac{L}{\Gamma(\alpha - p + 1)} \|u - v\|. \tag{3.17}$$

Thus from (3.16) and (3.17), we have

$$\|Tu - Tv\| \leq \left(\frac{2L}{\Gamma(\alpha + 1)} + \frac{L}{\Gamma(\alpha - p + 1)} \right) \|u - v\| = L\Omega \|u - v\|.$$

Thus by applying Theorem 2.6, T has a unique fixed point which is the unique positive solution of BVP (1.1). □

4. EXAMPLE

Example 1.

$$\begin{cases} \mathcal{D}^{\frac{5}{2}} u(t) = \frac{t}{4} + \frac{\sin^{\frac{1}{2}} |u(t)|}{32} + \frac{\cos |\mathcal{D}^{\frac{1}{2}} u(t)|}{32}, t \in [0, 1], \\ \mathcal{I}^{\frac{1}{2}} u(0) = \mathcal{D}^{\frac{1}{2}} u(0) = u(1) = 0. \end{cases} \tag{4.1}$$

From (4.1), we see that $\alpha = \frac{5}{2}$ and $p = \frac{1}{2}$, where

$$f(t, u, \mathcal{D}^p u) = \frac{t}{4} + \frac{\sin^{\frac{1}{2}} |u(t)|}{32} + \frac{\cos^{\frac{1}{2}} |\mathcal{D}^{\frac{1}{2}} u(t)|}{32}$$

$$|f(t, u, \mathcal{D}^p u)| \leq \frac{1}{4} + \frac{1}{32} |u|^{\frac{1}{2}} + \frac{1}{32} |\mathcal{D}^{\frac{1}{2}} u(t)|^{\frac{1}{2}},$$

clearly $a_0 = \frac{1}{4}$, $b = \frac{1}{32}$, $c = \frac{1}{32}$, $\theta = \rho = \frac{1}{2}$, $u, v \in R$, then

$$|f(t, u, \mathcal{D}^p u) - f(t, v, \mathcal{D}^p v)| \leq \frac{1}{32} (|u - v| + |\mathcal{D}^p u - \mathcal{D}^p v|), \text{ where } L = \frac{1}{32}.$$

As assumptions (A_1) to (A_3) hold, so BVP (4.1) has at least one solution. While

$$\Omega = \frac{2}{\Gamma(\frac{5}{2} + 1)} + \frac{1}{\Gamma(\frac{5}{2} - \frac{1}{2} + 1)} = \frac{2}{\Gamma(\frac{7}{2})} + \frac{1}{\Gamma(3)} = \frac{2}{\frac{5}{2} \cdot \frac{3}{2} \cdot \frac{1}{2} \Gamma(\pi)} + \frac{1}{2} = 1.10180.$$

From which, we have $L\Omega = \frac{1.10180}{32} = 0.034431 < 1$. Hence BVP (4.1) has a unique solution.



5. CONCLUSION

By the use of classical fixed point theorem of nonlinear alternative of Leray-Schauder type and Banach concatenation theorem, we have developed some necessary and sufficient conditions for the existence of at least one positive solutions to a class of nonlinear fractional differential equations. Where the nonlinearity of the problem explicitly depends on the fractional order derivative which produce an extra complexity and such type of problems occur in applications in real world.

ACKNOWLEDGMENT

We are really thankful to the anonymous referee and editorial board for their valuable suggestions, corrections and comments, which improved the quality of this paper.

REFERENCES

- [1] R. P. Agarwal, M. Benchohra and S. Hamani, Boundary value problems for differential inclusions with fractional order, *Adv. Stud. Contemp. Math*, 12(2) (2008), 181–196.
- [2] B. Ahmad and J. J. Nieto, Existence results for nonlinear boundary value problems of fractional integrodifferential equations with integral boundary conditions, *Bound. Value Probl. Art*, (2009) Article ID 708576 (2009), 11.
- [3] A.Ali and R.A.Khan, Existence of Solutions of Fractional Differential Equations via Topological Degree Theory, *J.Comput. Theo. Nanoscience*, 13 (2016), 1–5,
- [4] Z. Bai, On positive solutions of a nonlocal fractional boundary value problem, *Nonlinear Anal: Theo. Meth. Appl*, 72(2) (2010), 916–924.
- [5] C.Z.Bai, Triple positive solutions for a boundary value problem of nonlinear fractional differential equations, *Elect. J. Qual. Theo. Diff. Equns*, 24 (2008), 1–10.
- [6] A. Belarbi, M. Benchohra and A. Ouahab, Existence results for functional differential equations of fractional order, *Appl. Anal*, 85 (2006), 1459–1470.
- [7] M. Benchohra, J. Henderson, S. K. Ntouyas, and A. Ouahab, Existence results for fractional order functional differential equations with infinite delay, *J. Math. Anal. Appl*, 332 (2008), 1340–1350.
- [8] M. Benchohra, S. Hamani and S. K. Ntouyas, Boundary value problems for differential equations with fractional order and nonlocal conditions, *Nonlinear. Anal: Theo. Meth. Appl*, 71 (2009), 2391-2396.
- [9] M. Caputo, Linear Models of dissipation whose Q is almost frequency independent, *Int. J. Geo. Sci*, 13(5) (1967), 529–539.
- [10] Z. Cui, P. Yu and Z. Mao, Existence of solutions for nonlocal boundary value problems of nonlinear fractional differential equations, *Advanc. Dyn. Sys. Appl*, 7(1) (2012), 243–254.
- [11] K.S. Fa, Generalized Langevin equation with fractional derivative and long-time correlation function, *Phys. Rev.E*, 73 (2006), Article ID 061104, 3345–3356.
- [12] C. Goodrich, Existence of positive solution to a class of fractional differential equations, *Appl. Math. Lett*, 23 (2010), 1050–1055.
- [13] K. Ghanbari, Y. Gholami, Existence and multiplicity of positive solutions for a coupled system of perturbed nonlinear fractional differential equations, *Comput. Method. Diff. Equns*, 1(1) 2013, 39–54
- [14] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
- [15] A. A.Kilbas, H. M. Srivastava, and J. J Trujillo, *Theory and Applications of Fractional Differential Equation*, North-Holland arthematics Studies, Elsevier, Amsterdam, 2006.
- [16] N. Kosmatov, A singular boundary value problems nonlinear differential equations of fractional order, *J. Appl. Math. Comput*, 29(1) (2009), 125–135.



- [17] V. Lakshmikantham, S. Leela and J. Vasundhara, *Theory of Fractional Dynamic Systems*, Cambridge Academic Publishers, Cambridge, UK, 2009.
- [18] C. F. Li, X. N. Luo and Y. Zhou, Existence of positive solutions of the boundary value problem for nonlinear fractional differential equations, *Comput. Math. Appl*, 59(3) (2010), 1363–1375.
- [19] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [20] A. M. A. EL-Sayed and E.O.Bin-Taher, Positive solutions for a nonlocal multi-point boundary-value problem of fractional and second order, *Electr. J. Diff. Equns*, 64 (2013), 1–8.
- [21] K.Shah, S. Zeb and R. A.Khan, Existence and uniqueness of solution of fractional order m-point boundary value problems, *Frac. Diff. Calc*, 5(2) (2015), 171–181.
- [22] K.Shah, A.Ali and R.A.Khan, Degree theory and existence of positive solutions to coupled systems of multi-point boundary value problems, *Boundary Value Problems*, 2016(43) (2016), 1–12.
- [23] K.Shah and R.A.Khan, Iterative scheme to a coupled system of highly nonlinear fractional order differential equations, *Comput. Method. Differ. Equns*, 3(3) (2015), 163–176.
- [24] X. Su and S. Zhang, Unbounded solutions to a boundary value problem of fractional order on the half-line, *Comput. Math. Appl*, 61 (2011), 1079–1087.
- [25] S. Picozzi, B. West, Fractional Langevin model of memory in financial markets, *Phys. Rev. E*, 66 (2002), Article ID 046118, 1023–1032.
- [26] I. Podlubny, *Fractional Differential Equations*, Mathematics in Science and Engineering, Academic Press, New York, 1999.
- [27] E. Zeidler, *Nonlinear Functional Analysis and its Applications*, part II/B: Nonlinear Monotone Operators, Springer Verlag, New York, 1989.
- [28] W. Zhong and W. Lin, Nonlocal and multiple-point boundary value problem for fractional differential equations, *Comput. Math. Appl*, 59(3) (2010), 1345–1351.

