



## Plasma particles dispersion based on Bogoyavlensky-Konopelchenko mathematical model

Ahmed S. Rashed<sup>1,2</sup>, Mokhtar Mohamed<sup>2</sup>, and Mustafa Inc<sup>3,\*</sup>

<sup>1</sup>Department of Physics and Engineering Mathematics Department, Faculty of Engineering, Zagazig University, Egypt.

<sup>2</sup>Faculty of Engineering, Delta University for Science and Technology, Gamasa, Egypt.

<sup>3</sup>Department of Mathematics, Firat University, 23119 Elazig, Turkiye.

### Abstract

An optimal system of Lie infinitesimals has been used in an investigation to find a solution to the (2+1)-dimensional Bogoyavlensky-Konopelchenko equation (BKE). This investigation was conducted to characterize certain fantastic characteristics of plasma-particle dispersion. A careful investigation into the Lie space with an unlimited number of dimensions was carried out to locate the relevant arbitrary functions. When developing accurate solutions for the BKE, it was necessary to establish an optimum system that could be employed in single, double, and triple combination forms. There were some fantastic wave solutions developed, and these were depicted visually. The Optimal Lie system demonstrates that it can obtain many accurate solutions to evolution equations.

**Keywords.** Evolution equations, BKE, Lie infinitesimals, Optimal Lie system.

**2010 Mathematics Subject Classification.** 37L05, 17B45.

### 1. INTRODUCTION

Investigating nonlinear physical processes plays a crucial role in understanding various scientific and technological phenomena. A key aspect of this investigation is the analysis of precise traveling wave solutions to nonlinear partial differential equations. These solutions provide valuable insights into the behavior of nonlinear waves and are essential for advancing knowledge in many fields, particularly in industry. Nonlinear wave phenomena are prevalent across diverse disciplines, spanning fluid mechanics, plasma physics, optical fibers, biology, solid-state physics, chemical kinematics, chemical physics, and geochemistry, among others. By studying the intricate dynamics of nonlinear waves in these areas, researchers can uncover fundamental principles and develop practical applications. One widely recognized evolution equation used in various contexts is the Bogoyavlensky-Konopelchenko equation (BKE). This equation's versatility allows it to be applied in different settings, making it a valuable tool for investigating nonlinear wave phenomena. For instance, the BKE can help explain the dispersion of particles in plasma, which occurs due to the diffusion of plasma. By employing this theory, scientists can gain a better understanding of plasma dynamics and its implications in fields such as fusion energy research and astrophysics. Numerous methods have been employed to attain exact solutions of these equations, such as Backlund transform [1–4], auxiliary function method [5], extended tanh method [6–9], Lie group method [10–15], Darboux transformation [16–19], the projective Riccati equations method [20, 21], sine-cosine method [9, 22, 23] and the variation iteration method [24–26]. Moreover, sub-equation method [27, 28], nonlocal conservation laws [29], Painlevé analysis via WTCKruskal algorithm [30] are also some useful methods. Similarity transformation methods, which can be used to construct either exact solutions or reduce the partial differential equation (PDE) model into an ordinary differential equation (ODE) model, are one of the most powerful techniques. These methods include Lie infinitesimals, group transformation methods, and hidden symmetries, among others, and can be used to construct either exact solutions or reduce the PDE model into an ODE model [31–42]. The motivation of this manuscript is to discover new families of solutions compatible with describing the dispersion

Received: 06 January 2023 ; Accepted: 09 February 2024.

\* Corresponding author. Email: minc@firat.edu.tr .

of plasma particles. The technique is to investigate and create an optimal system of Lie space instead of using the typical Lie infinitesimals directly. Moreover, the hidden symmetry vectors are detected during the reduction process, which will lead to totally new families of solutions. The paper is arranged as follows. In section 2, the mathematical formulation of the problem is introduced with Lie infinitesimals vectors. Then, the commutator and adjoint tables are constructed in section 3. The optimal system is discovered in section 4, and the results are illustrated in section 5. Finally, the paper is terminated by conclusion remarks.

## 2. MATHEMATICAL FORMULATION

BKE in (2+1) dimensions may describe the dispersion of plasma particles. One of its forms can be written as:

$$u_{xt} + u_{xxx}y + 4u_xu_{xy} + 2u_{xx}u_y = 0, \quad (2.1)$$

$$\begin{aligned} X_1 &= \frac{\partial}{\partial t} + F_1(t)\frac{\partial}{\partial x} + \left(\frac{y}{2}\frac{d}{dt}F_1(t) + F_2(t)\right)\frac{\partial}{\partial u}, \\ X_2 &= F_3(t)\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \left(\frac{y}{2}\frac{d}{dt}F_3(t) + F_4(t)\right)\frac{\partial}{\partial u}, \\ X_3 &= F_5(t)\frac{\partial}{\partial x} + t\frac{\partial}{\partial y} + \left(\frac{y}{2}\frac{d}{dt}F_5(t) + F_6(t) + \frac{x}{4}\right)\frac{\partial}{\partial u}, \\ X_4 &= t\frac{\partial}{\partial t} + \left(F_7(t) + \frac{x}{2}\right)\frac{\partial}{\partial x} + \left(\frac{y}{2}\frac{d}{dt}F_7(t) + F_8(t) - \frac{u}{2}\right)\frac{\partial}{\partial u}, \\ X_5 &= \left(F_9(t) - \frac{x}{2}\right)\frac{\partial}{\partial x} + y\frac{\partial}{\partial y} + y\frac{\partial}{\partial y} + \left(\frac{y}{2}\frac{d}{dt}F_9(t) + F_{10}(t) + \frac{u}{2}\right)\frac{\partial}{\partial u}, \\ X_6 &= \frac{t^2}{2}\frac{\partial}{\partial t} + \left(F_{11}(t) + \frac{xt}{4}\right)\frac{\partial}{\partial x} + \frac{yt}{2}\frac{\partial}{\partial y} + \left(\frac{y}{2}\frac{d}{dt}F_{11}(t) + F_{12}(t) - \frac{ut}{4} + \frac{xy}{8}\right)\frac{\partial}{\partial u}. \end{aligned} \quad (2.2)$$

Where,  $F_i(t)$ ,  $i = 1..12$  are arbitrary functions to be determined later.

## 3. COMMUTATOR AND ADJOINT TABLES

To evaluate the appropriate functions, the commutator table of the vectors  $X_1, X_2, X_3, X_4, X_5$  and  $X_6$  is constructed using the relation  $[X_i, X_j] = X_i(X_j) - X_j(X_i)$ . For example, the following commutators are computed.

$$\begin{aligned} [X_1, X_2] &= \left(\frac{d}{dt}F_3(t)\right)\frac{\partial}{\partial x} + \left(\frac{y}{2}\frac{d^2}{dt^2}F_3(t) + \frac{d}{dt}F_4(t) - \frac{1}{2}\frac{d}{dt}F_1(t)\right), \\ [X_1, X_3] &= \left(\frac{d}{dt}F_5(t)\right)\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \left(\frac{y}{2}\frac{d^2}{dt^2}F_5(t) + \frac{d}{dt}F_6(t) + \frac{1}{4}F_1(t) - \frac{t}{2}\frac{d}{dt}F_1(t)\right)\frac{\partial}{\partial u}, \\ [X_1, X_4] &= \frac{\partial}{\partial t} + \left(\frac{d}{dt}F_7(t) + \frac{1}{2}F_1(t) - t\frac{d}{dt}F_1(t)\right)\frac{\partial}{\partial x} \\ &+ \left(\frac{y}{2}\frac{d^2}{dt^2}F_7(t) + \frac{d}{dt}F_8(t) - \frac{y}{4}\frac{d}{dt}F_1(t) - \frac{1}{2}F_2(t) - \frac{ty}{2}\frac{d}{dt}F_1(t) - t\frac{d}{dt}F_2(t)\right)\frac{\partial}{\partial u}, \\ [X_1, X_5] &= \left(\frac{d}{dt}F_9(t) - \frac{1}{2}F_1(t)\right)\frac{\partial}{\partial x} + \left(\frac{y}{2}\frac{d^2}{dt^2}F_9(t) + \frac{d}{dt}F_{10}(t) + \frac{y}{4}\frac{d}{dt}F_1(t) + \frac{1}{2}F_2(t) - \frac{y}{2}\frac{d}{dt}F_1(t)\right)\frac{\partial}{\partial u}, \end{aligned}$$



$$\begin{aligned}
 [X_1, X_6] &= t \frac{\partial}{\partial t} + \left( \frac{d}{dt} F_{11}(t) + \frac{x}{4} + \frac{t}{4} F_1(t) - \frac{t^2}{2} \frac{d}{dt} F_1(t) \right) \frac{\partial}{\partial x} + \frac{y}{2} \frac{\partial}{\partial y} \\
 &+ \left( \frac{y}{2} \frac{d^2}{dt^2} F_{11}(t) + \frac{d}{dt} F_{12}(t) - \frac{u}{4} + \frac{y}{8} F_1(t) - \frac{ty}{8} \frac{d}{dt} F_1(t) - \frac{t}{4} F_2(t) - \frac{t^2 y}{4} \frac{d^2}{dt^2} F_1(t) \right) \frac{\partial}{\partial u} \\
 &\left( -\frac{t^2}{2} \frac{d}{dt} F_2(t) - \frac{yt}{4} \frac{d}{dt} F_1(t) \right) \frac{\partial}{\partial u}.
 \end{aligned} \tag{3.1}$$

To get a solvable system, the following condition must be satisfied.  $[X_i, X_j] = \lambda X_n$  where  $n = 1 \dots 6$ .

Setting  $[X_1, X_2] = 0$  leads to:

$$F_3(t) = 0 \quad \text{and} \quad F_4(t) = \frac{1}{2} F_1(t). \tag{3.2}$$

Setting  $[X_1, X_3] = X_2$  leads to:

$$F_5(t) = 0, \quad F_4(t) = \frac{1}{2} F_1(t) \quad \text{and} \quad \frac{d}{dt} F_6(t) = \frac{1}{4} F_1(t) + \frac{t}{2} \frac{d}{dt} F_1(t). \tag{3.3}$$

Moreover, setting  $[X_1, X_4] = X_1$  leads to:

$$\frac{d}{dt} F_7(t) = \frac{1}{2} F_1(t) + t \frac{d}{dt} F_1(t), \tag{3.4}$$

$$\frac{y}{2} \frac{d^2 F_7}{dt^2} + \frac{d F_8}{dt} - \frac{y}{4} \frac{d F_1}{dt} - \frac{F_2}{2} - \frac{ty}{2} \frac{d^2 F_1}{dt^2} - t \frac{d F_2}{dt} = \frac{y}{2} \frac{d F_1}{dt} + F_2. \tag{3.5}$$

Also, for  $[X_1, X_5] = 0$ , one can get

$$\frac{d F_9}{dt} - \frac{F_1}{2} = 0, \quad \frac{y}{2} \frac{d^2 F_9}{dt^2} + \frac{d F_{10}}{dt} - \frac{y}{4} \frac{d F_1}{dt} + \frac{F_2}{2} = 0. \tag{3.6}$$

In the same way, the remaining commutators are simplified, and the resultant ordinary differential equations are solved simultaneously to get:

$$\begin{aligned}
 F_1 &= 2, \quad F_2 = 2, \quad F_3 = 0, \quad F_4 = 1, \quad F_5 = 0, \quad F_6 = \frac{t}{2}, \quad F_7 = t, \quad F_8 = 3t, \quad F_9 = t, \quad F_{10} = -t, \\
 F_{11} &= \frac{t^2}{2}, \quad F_{12} = \frac{3}{2} t^2.
 \end{aligned} \tag{3.7}$$

Thus, the space of Lie vectors will be:

$$\begin{aligned}
 X_1 &= \frac{\partial}{\partial t} + 2 \frac{\partial}{\partial x} + 2 \frac{\partial}{\partial u}, \\
 X_2 &= \frac{\partial}{\partial y} + \frac{\partial}{\partial u}, \\
 X_3 &= t \frac{\partial}{\partial y} + \left( \frac{t}{2} + \frac{x}{4} \right) \frac{\partial}{\partial u}, \\
 X_4 &= t \frac{\partial}{\partial t} + \left( t + \frac{x}{2} \right) \frac{\partial}{\partial x} + \left( \frac{y}{2} + 3t - \frac{u}{2} \right) \frac{\partial}{\partial u}, \\
 X_5 &= \left( t - \frac{x}{2} \right) \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + \left( \frac{y}{2} - t + \frac{u}{2} \right) \frac{\partial}{\partial u},
 \end{aligned}$$



$$X_6 = \frac{t^2}{2} \frac{\partial}{\partial t} + \left( \frac{t^2}{2} + \frac{xt}{4} \right) \frac{\partial}{\partial x} + \frac{yt}{2} \frac{\partial}{\partial y} + \left( \frac{yt}{2} + \frac{3}{2}t^2 - \frac{ut}{4} + \frac{xy}{8} \right) \frac{\partial}{\partial u}. \quad (3.8)$$

Finally, the commutator table is constructed hereafter:

TABLE 1. Commutator Table.

$[X_i, X_j]$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6$
$\mathbf{X}_1$	0	0	$X_2$	$X_1$	0	$X_4 + \frac{1}{2}X_5$
$\mathbf{X}_2$	0	0	0	0	$X_2$	$\frac{1}{2}X_3$
$\mathbf{X}_3$	$-X_2$	0	0	$-X_3$	$X_3$	0
$\mathbf{X}_4$	$-X_1$	0	$X_3$	0	0	$X_6$
$\mathbf{X}_5$	0	$-X_2$	$-X_3$	0	0	0
$\mathbf{X}_6$	$-X_4 - \frac{1}{2}X_5$	$\frac{1}{2}X_3$	0	$-X_6$	0	0

The adjoint table is to be constructed using the relation:

$$\text{Ad}(e^{\varepsilon V}) W_0 = W_0 - \varepsilon [V, W_0] + \frac{\varepsilon^2}{2} [V, [V, W_0]] - \dots \quad (3.9)$$

Using this relation, the adjoint table is constructed in the form:

TABLE 2. Adjoint Table.

Ad	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6$
$\mathbf{X}_1$	$X_1$	$X_2$	$X_3 - \varepsilon X_2$	$X_4 - \varepsilon X_1$	$X_5$	$X_6 - \varepsilon \left( X_4 + \frac{1}{2}X_5 \right) + \frac{\varepsilon^2}{2} X_1$
$\mathbf{X}_2$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5 - \varepsilon X_2$	$X_6 - \frac{\varepsilon}{2} X_3$
$\mathbf{X}_3$	$X_1 + \varepsilon X_2$	$X_2$	$X_3$	$X_4 + \varepsilon X_3$	$X_5 - \varepsilon X_3$	$X_6$
$\mathbf{X}_4$	$e^{-\varepsilon} X_1$	$X_2$	$e^{-\varepsilon} X_3$	$x_4$	$X_5$	$e^{-\varepsilon} X_6$
$\mathbf{X}_5$	$X_1$	$e^{\varepsilon} X_2$	$e^{\varepsilon} X_3$	$X_4$	$X_5$	$X_6$
$\mathbf{X}_6$	$X_1 + \varepsilon \left( X_4 + \frac{1}{2}X_5 \right) + \frac{\varepsilon^2}{2} X_6$	$X_2 + \varepsilon X_3$	$X_3$	$X_4 + \varepsilon X_6$	$X_5$	$X_6$

#### 4. OPTIMAL LIE SYSTEM

The following adjoint matrices are constructed from Table 2.

$$\text{Ad}(e^{\varepsilon_1 X_1}) = \begin{bmatrix} 1 & 0 & 0 & -\varepsilon_1 & 0 & \frac{\varepsilon_1^2}{2} \\ 0 & 1 & -\varepsilon_1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & -\varepsilon_1 \\ 0 & 0 & 0 & 0 & 1 & \frac{-\varepsilon_1}{2} \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$\text{Ad}(e^{\varepsilon_2 X_2}) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & -\varepsilon_2 & 0 \\ 0 & 0 & 1 & 0 & 0 & \frac{-\varepsilon_2}{2} \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$



$$\text{Ad}(e^{\varepsilon_3 X_3}) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ \varepsilon_3 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & \varepsilon_3 & -\varepsilon_3 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$\text{Ad}(e^{\varepsilon_4 X_4}) = \begin{bmatrix} e^{-\varepsilon_4} & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{-\varepsilon_4} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & e^{-\varepsilon_4} \end{bmatrix},$$

$$\text{Ad}(e^{\varepsilon_5 X_5}) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & e^{-\varepsilon_5} & 0 & 0 & 0 & 0 \\ 0 & 0 & e^{-\varepsilon_5} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

$$\text{Ad}(e^{\varepsilon_6 X_6}) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & \varepsilon_6 & 1 & 0 & 0 & 0 \\ \varepsilon_6 & 0 & 0 & 1 & 0 & 0 \\ \frac{\varepsilon_6}{2} & 0 & 0 & 0 & 1 & 0 \\ \frac{\varepsilon_6}{2} & 0 & 0 & 0 & 0 & 1 \end{bmatrix}. \tag{4.1}$$

Now  $Ad_g = Ad(e^{\varepsilon_1 X_1}) * Ad(e^{\varepsilon_2 X_2}) * \dots * Ad(e^{\varepsilon_6 X_6})$ .  
 Let:

$$\frac{1}{a} Ad_g \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \\ \alpha_5 \\ \alpha_6 \end{bmatrix} = \begin{bmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \\ \beta_4 \\ \beta_5 \\ \beta_6 \end{bmatrix}. \tag{4.2}$$

To obtain the optimal system, the following cases are studied.

**Case 1:**

For  $\alpha_6 \neq 0$ , there are two subcases:

If  $\varepsilon_1 = 0$ , the vector  $X_3$  is an optimal vector,

If  $\varepsilon_2 = 0$ , the vector  $X_1 + X_4 + X_5$  is an optimal vector.

**Case 2:**

For  $\alpha_5 \neq 0$ , there are three subcases:



If  $\varepsilon_3 \neq 0$ , the vector  $X_2 + X_3 + X_5$  is an optimal vector,

If  $\varepsilon_3 = 0$ , the vector  $X_2 + X_5$  is an optimal vector,

If  $\varepsilon_2 = \varepsilon_3 = 0$ , the vector  $X_5$  is an optimal vector.

**Case 3:**

For  $\alpha_4 \neq 0$ , the vector  $X_4$  is an optimal vector.

**Case 4:**

For  $\alpha_3 \neq 0$ , the vector  $X_3$  is an optimal vector.

**Case 5:**

For  $\alpha_2 \neq 0$ , the vector  $X_2$  is an optimal vector.

**Case 6:**

For  $\alpha_1 \neq 0$ , the vector  $X_1$  is an optimal vector.

**Case 7:**

For  $\alpha_4 \neq 0, \alpha_5 \neq 0$ , the vector  $X_1 + X_4$  is an optimal vector.

Following the same procedures, all cases were studied and the optimal system of Lie vectors are:

**Single Vectors:**  $X_1, X_2, X_3, X_4, X_5$ .

**Double linear combinations:**  $X_1 + X_4, X_2 + X_4, X_1 + X_3, X_4 + X_5, X_1 + X_5$ .

**Triple linear combinations:**  $X_1 + X_4 + X_5, X_2 + X_3 + X_5$ .

These vectors are used to reduce the Bogoyavlensky-Konopelchenko equation to an invariant ordinary differential equation to get the exact solutions.

## 5. RESULTS AND DISCUSSION

5.1. **Using  $X_1$ .** The Equation (2.1) is reduced to:

$$w_{rrrs} + (2w_s - 2) + 4w_r w_{rs} = 0, \quad (5.1)$$

where,  $r = -2t + x, s = y, w(r, s) = u(t, x, y) - 2t$ .

Equation (5.1) has the following exact solution:

$$w(r, s) = 2C_2 \tanh \left( C_2 r + \frac{s}{2C_2} + C_1 \right) + C_3. \quad (5.2)$$

Then, by back substitution, the solution of Equation (2.1) is:

$$u(x, y, t) = 2t + 2C_2 \tanh \left( C_2(x - 2t) + \frac{y}{2C_2} + C_1 \right) + C_3. \quad (5.3)$$

Moreover, when studying the symmetry vectors of (5.1), these vectors can be used to reduce the equation to an ODE.

Using the vector  $\frac{\partial}{\partial r} + \frac{\partial}{\partial s} + \frac{\partial}{\partial w}$ , Equation (5.1) is reduced to:

$$\frac{d^4 V(\alpha)}{d\alpha^4} - \left( 6 \frac{dV(\alpha)}{d\alpha} - 6 \right) \frac{d^2 V(\alpha)}{d\alpha^2} = 0, \quad (5.4)$$

where,  $\alpha = -r + s, V(\alpha) = w(r, s) - r$ .

Now, one of the symmetry vectors of (5.4) is:

$\frac{\partial}{\partial V}$  which reduces its order to:

$$\psi''' = (6\psi - 6)\psi', \text{ where, } \varepsilon = \alpha, \psi(\alpha) = \frac{dV(\alpha)}{d\alpha}. \quad (5.5)$$

Moreover, this Equation, (5.5), has one of its vectors as  $\frac{\partial}{\partial \varepsilon}$  which reduces the order of (5.5) to:

$$TT'' = -T''^2 + 6\delta - 6, \quad (5.6)$$

where

$$\delta = \psi(\varepsilon), T(\delta) = \frac{d\psi(\varepsilon)}{d\varepsilon},$$



which has a solution in the form:

$$T(\delta) = \pm\sqrt{2\delta^3 - 6\delta^2 - C_1\delta + C_2}. \tag{5.7}$$

Then, by back substitution, the solution of Equation (1) is written as:

$$u(x, y, t) = x + \sqrt{6} \tan\left(\frac{\sqrt{6}}{2} (y + 2t - x) + C\right). \tag{5.8}$$

5.2. **Using  $X_2$ .** By the same procedures,  $X_2$  is used to get the following solution:

$$u(x, y, t) = y + F_1(t) + F_2(-2t + x), \tag{5.9}$$

where,  $F_1$  and  $F_2$  are arbitrary functions in their arguments.

5.3. **Using  $X_3$ .** By the same procedures,  $X_3$  is used to attain the following solution:

$$u(x, y, t) = \frac{(2t + x)y}{4t} + F_2(t) + \left(\frac{1}{\sqrt{t}}\right) F_1\left(\frac{x - 2t}{\sqrt{t}}\right), \tag{5.10}$$

where,  $F_1$  and  $F_2$  are arbitrary functions in their arguments.

Using the other optimal vectors, one can get the following solutions:

$$u(x, y, t) = t\left(2 + \frac{x}{4}\right) + C_2\left(-\frac{t^2}{2} + y\right) - \frac{(-2t + x)^2}{8(2C_2 - 2)} + C_3(-2t + x) + C_4, \tag{5.11}$$

$$u(x, y, t) = \frac{(2t + x + 4)y}{4t + 4} + F_2(t) + \left(\frac{1}{\sqrt{t + 1}}\right) F_1\left(\frac{x - 2t}{\sqrt{t + 1}}\right), \tag{5.12}$$

$$u(x, y, t) = 3t + 2C_2 \tanh\left(C_2(y - 2t) + 2C_2\frac{y - t}{4C_2^2 - 1} + C_1\right) + C_3, \tag{5.13}$$

Some of the previous results are depicted in Figures 1-4. These figures illustrate some interpretation of plasma particles dispersion in different cases of boundary conditions. These behaviors vary between kink solutions, trigonometric and arbitrary functions.




FIGURE 1. Solution behavior of Equation (5.3).






FIGURE 2. Solution behavior of Equation (5.8).




FIGURE 3. Solution behavior of Equation (5.9) with  $F_1(t) = \cos(t)$ ,  $F_2(x - 2t) = (x - 2t)e^{-(x - 2t)^2}$ .




FIGURE 4. Solution behavior of Equation (5.10) with  $F_2(t) = e^{-t}$ ,  $F_1\left(\frac{x-2t}{\sqrt{t}}\right) = 10 * \cos\left(\frac{x-2t}{\sqrt{t}}\right)$ .



The illustrations shown in Figures 1-4 show the behaviour of plasma particles in different cases. For example, in Figure 1, the graph shows the case of limited dispersion of the particles. In Figure 2, the periodic behavior of the plasma particles is illustrated. Some chaotic arbitrary dispersion of the particles are shown in Figure 3 and 4 due to the arbitrary functions in the solution.

## 6. CONCLUSION

The solution of the (2+1)-dimensional BKE has been investigated using an optimal system of Lie infinitesimals to describe some fabulous behaviors of plasma-particles dispersion. The infinite-dimensional Lie space was precisely analyzed to detect the appropriate arbitrary functions. An optimal system was obtained as single, double, and triple combinations to be used in constructing exact solutions of the BKE. Some fabulous wave solutions were constructed and illustrated graphically. The Optimal Lie system proves its ability to get numerous exact solutions to evolution equations.

## REFERENCES

- [1] S. Yadong, *Backlund transformation, lax pairs and explicit exact solutions for the shallow water waves equation*, Applied Mathematics and Computation, *187* (2007), 1286-1297.
- [2] Z. L. J. Su, and F. Xie, *Construction of exact solutions to the jimboniwa equation through bcklund transformation and symbolic computation*, Computers & Mathematics with Applications, *65* (2013), 648-565.
- [3] B. Lu, *Coupling bcklund trasnsformation of riccati equation and division theorem method for traveling wave solutions of some class of NLPDEs*, Communications in Nonlinear Science and Numerical Simulation, *17* (2012), 4626-4633.
- [4] J. M. Conde, P. R. Gordo, and A. Pickering, *A new kind of bcklund transformation for partial differential equations*, Reports on Mathematical Physics, *70* (2012), 149-161.
- [5] Y. Feng and H. Q. Zhang, *A new auxiliary function method for solving the generalized coupled hirotasatsuma kdv system*, Applied Mathematics and Computation, *200* (2008), 283-288.
- [6] A. Bekir, *Applications of the extended tanh method for coupled nonlinear evolution equations*, Communications in Nonlinear Science and Numerical Simulation, *13* (2008), 1748-1757.
- [7] S. Dubey and S. Chakraverty, *Application of modified extended tanh method in solving fractional order coupled wave equations*, Mathematics and Computers in Simulation, *198* (2022), 509-520.
- [8] A. A. Mamun, S. N. Ananna, P. P. Gharami, T. An, and M. Asaduzzaman, *The improved modified extended tanh-function method to develop the exact travelling wave solutions of a family of 3D fractional WBBM equations*, Results in Physics, *41* (2022), 105969.
- [9] M. Ali, M. Alquran, and O. B. Salman, *A variety of new periodic solutions to the damped (2+1)-dimensional Schrodinger equation via the novel modified rational sine-cosine functions and the extended tanh-coth expansion methods*, Results in Physics, *37* (2022), 105462.
- [10] H. A. Zedan, *Blow-up the symmetry analysis for the hirota-satsuma equations*, Journal of Contemporary Mathematical Analysis, *43* (2008), 108-117.
- [11] K. Singh and R. K. Gupta, *Lie symmetries and exact solutions of a new generalized hirotasatsuma coupled KdV system with variable coefficients*, International Journal of Engineering Science, *44* (2006), 241-255.
- [12] A. S. Rashed and M. M. Kassem, *Hidden symmetries and exact solutions of integro-differential Jaulent-Miodek evolution equation*, Applied Mathematics and Computation, *247* (2014), 1141-1155.
- [13] S. M. Mabrouk and A. S. Rashed, *Analysis of (3 + 1)-dimensional Boiti-Leon-Manna-Pempinelli equation via lax pair investigation and group transformation method*, Computers & Mathematics with Applications, *74* (2017), 2546-2556.
- [14] M. M. Kassem and A. S. Rashed, *N-solitons and cuspon waves solutions of (2+1)-dimensional Broer-Kaup-Kupershmidt equations via hidden symmetries of Lie optimal system*, Chinese Journal of Physics, *57* (2019), 90-104.



- [15] R. Saleh and A. S. Rashed, *New exact solutions of  $(3 + 1)$ -dimensional generalized Kadomtsev-Petviashvili equation using a combination of lie symmetry and singular manifold methods*, *Mathematical Methods in the Applied Sciences*, *43* (2020), 2045-2055.
- [16] X. L. Yong and H. Q. Zhang, *New exact solutions to the generalized coupled Hirota-Satsuma KdV system*, *Chaos, Solitons & Fractals*, *26* (2005), 1105-1110.
- [17] L. Guo, Y. Cheng, D. Mihalache, and J. He, *Darboux transformation and higher-order solutions of the Sasatsuma equation*, *Romanian Journal of Physics*, *64* (2019), 104.
- [18] F. C. Fan and Z.G. Xu, *Breather and rogue wave solutions for the generalized discrete Hirota equation via Darboux-Bäcklund transformation*, *Wave Motion*, *119* (2023), 103139.
- [19] W. K. Xie and F.C. Fan, *Soliton, breather, rogue wave and continuum limit in the discrete complex modified Korteweg-de Vries equation by Darboux-Bäcklund transformation*, *Journal of Mathematical Analysis and Applications*, *525* (2023), 127251.
- [20] H. C. Hu and Q. P. Liu, *New Darboux transformation for Hirota-Satsuma coupled KdV system*, *Chaos, Solitons & Fractals*, *17* (2003), 921-928.
- [21] D. Lu, B. Hong, and L. Tian, *New explicit exact solutions for the generalized coupled Hirota-Satsuma KdV system*, *Computers and Mathematics with Applications*, *53* (2007), 1181-1190.
- [22] S. A. El-Wakil, M. A. Abdou, and A. Hendi, *New periodic and soliton solutions of nonlinear evolution equations*, *Applied Mathematics and Computation*, *197* (2008), 497-506.
- [23] S. W. Yao, S. Behera, M. Inc, H. Rezazadeh, J. P. S. Virdi, W. Mahmoud, O. Abu Arqub, and M. S. Osman, *Analytical solutions of conformable Drinfeld-Sokolov-Wilson and Boiti Leon Pempinelli equations via sine-cosine method*, *Results in Physics*, *42* (2022), 105990.
- [24] A. A. Soliman and M. A. Abdou, *Numerical solutions of nonlinear evolution equations using variational iteration method*, *Journal of Computational and Applied Mathematics*, *207* (2007), 111-120.
- [25] D. D. Ganji, M. Jannatabadi, and E. Mohseni, *Application of He's variational iteration method to nonlinear Jaulent-Miodek equations and comparing it with ADM*, *Journal of Computational and Applied Mathematics*, *207* (2007), 35-45.
- [26] Y. Nawaz, *Variational iteration method and homotopy perturbation method for fourth-order fractional integro-differential equations*, *Computers & Mathematics with Applications*, *61* (2011), 2330-2341.
- [27] H. Ypez-Martinez, A. Pashrashid, J. F. Gomez-Aguilar, L. Akinyemi, and H. Rezazadeh, *The novel soliton solutions for the conformable perturbed nonlinear Schrödinger equation*, *Modern Physics Letters B*, *36* (2022), 2150597.
- [28] L. Akinyemi, M. Enol, and O. S. Iyiola, *Exact solutions of the generalized multidimensional mathematical physics models via sub-equation method*, *Mathematics and Computers in Simulation*, *182* (2021), 211-233.
- [29] A. Souleymanou, A. Houwe, A. H. Kara, H. Rezazadeh, L. Akinyemi, S. P. T. Mukam, S. Y. Doka, and T. B. Bouetou, *Explicit exact solutions and conservation laws in a medium with competing weakly nonlocal nonlinearity and parabolic law nonlinearity*, *Optical and Quantum Electronics*, *55* (2023), 464.
- [30] L. Akinyemi and E. Morazara, *Integrability, multi-solitons, breathers, lumps and wave interactions for generalized extended Kadomtsev-Petviashvili equation*, *Nonlinear Dynamics*, *111* (2023), 4683-4707.
- [31] A. S. Rashed and M. M. Kassem, *Group analysis for natural convection from a vertical plate*, *Journal of Computational and Applied Mathematics*, *222* (2008), 392-403.
- [32] M. M. Kassem and A. S. Rashed, *Group solution of a time dependent chemical convective process*, *Applied Mathematics and Computation*, *215* (2009), 1671-1684.
- [33] S. M. Mabrouk and A. S. Rashed,  *$N$ -solitons, kink and periodic wave solutions for  $(3+1)$ -dimensional Hirota bilinear equation using three distinct techniques*, *Chinese Journal of Physics*, *60* (2019), 48-60.
- [34] A. S. Rashed, *Analysis of  $(3+1)$ -dimensional unsteady gas flow using optimal system of lie symmetries*, *Mathematics and Computers in Simulation*, *156* (2019), 327-346.
- [35] A. S. Rashed, S. M. Mabrouk, and A. M. Wazwaz, *Forward scattering for non-linear wave propagation in  $(3 + 1)$ -dimensional Jimbo-Miwa equation using singular manifold and group transformation methods*, *Waves in Random and Complex Media*, *32* (2022), 663-675.



- [36] A. S. Rashed, T. Mahmoud, and M. Kassem, *Analysis of homogeneous steady state nanofluid surrounding cylindrical solid pipes*, Egyptian Journal for Engineering Sciences and Technology, *31* (2020), 71-82.
- [37] A. S. Rashed, E. H. Nasr, and M. M. Kassem, *Similarity analysis of mass and heat transfer of fhd steady flow of nanofluid incorporating magnetite nanoparticles ( $fe_3o_4$ )*, East African Scholars Journal of Engineering and Computer Sciences, *3* (2020), 54-63.
- [38] A. S. Rashed, E. H. Nasr, and M. M. Kassem, *Boundary layer analysis adjacent to moving heated plate inside electrically conducting fluid with heat source/sink*, International Journal of Heat and Technology, *38* (2020), 682-688.
- [39] A. S. Rashed, S. M. Mabrouk, and A. M. Wazwaz, *Unsteady three-dimensional laminar flow over a submerged plate in electrically conducting fluid with applied magnetic field*, Waves in Random and Complex Media, (2021), 1-20.
- [40] A. S. Rashed, T. A. Mahmoud, and M. M. Kassem, *Behavior of nanofluid with variable brownian and thermal diffusion coefficients adjacent to a moving vertical plate*, Journal of Applied and Computational Mechanics, *7* (2021), 1466-1479.
- [41] A. S. Rashed, E. H. Nasr, and M. M. Kassem, *Mathematical investigation for flow characteristics of laminar ferro-nanofluid incorporating cobalt ferrite nanoparticles*, Journal of Nano Research, *68* (2021), 52-69.
- [42] R. Saleh, A. S. Rashed, and A. M. Wazwaz, *Plasma-waves evolution and propagation modeled by sixth order ramani and coupled ramani equations using symmetry methods*, Physica Scripta, *96* (2021), 085213.

Uncorrected Proof

