



## Application of a new method for nonlinear partial differential equations of fractional order arising in fluid mechanics

Oleg Kazakov<sup>1,\*</sup>, Nurgali Uteuliyev<sup>2</sup>, Diana Denisova<sup>3</sup>, Liya Shangaraeva<sup>4</sup>, and Galina Vukovich<sup>5</sup>

<sup>1</sup>Bryansk State University of Engineering and Technology, Russia.

<sup>2</sup>M. Auezov South Kazakhstan University, Shymkent, Kazakhstan.

<sup>3</sup>Russian State Social University, Moscow, Russia.

<sup>4</sup>Kazan Federal University, Kazan, Russia.

<sup>5</sup>Kuban State University, Krasnodar, Russia.

---

### Abstract

In this work, we established some exact solutions for the (2+1)-dimensional Zakharov-Kuznetsov, KdV, and K(2,2) equations which are considered based on the improved Exp-function method, by utilizing Maple software. We use the fractional derivatives with fractional complex transform. We obtained new periodic solitary wave solutions. The obtained solutions include three classes of soliton wave solutions in terms of hyperbolic function, trigonometric function, and rational function solutions. The obtained solutions and the exact solutions are shown graphically, highlighting the effects of non-linearity. Many other such types of nonlinear equations arising in fluid dynamics and nonlinear phenomena.

---

**Keywords.** Exp-function method(EFM), Zakharov-Kuznetsov, KdV, K(2,2).

**2010 Mathematics Subject Classification.** 02.60.Lj, 02.70.Wz, 02.90.+p.

### 1. INTRODUCTION

Many nonlinear physical phenomena arise in various fields of engineering and science such as fluid dynamics, nuclear reactor dynamics, plasma physics, biology, optical fibres and solid state physics. To describe these complex physical phenomena, nonlinear differential equations play a significant role. Therefore, obtaining the solutions of these nonlinear equations are a topic of great interest in the study of many fields of science. To better understand the workings of the physical problem, the mathematical model came into the picture in the form of nonlinear PDEs. The solutions of partial differential equations give the detailed summary about the nature of phenomena involved. Many numerical and analytical methods have been derived to deal with this kind of scientific problems. We need to adopt an effective and powerful method to investigate such type of mathematical model which gives the solutions upholding to physical reality. In most of the analytic techniques, linearization of the system is the main topic to focus on, and also, it is assumed that the nonlinearities are relatively insignificant. Sometimes, these assumptions made a strong affect on the solutions with respect to the real physics of the phenomena involved. Thus, finding the solutions of nonlinear ODEs and PDEs are still a significant problem. For this, we need new techniques to develop analytic and exact solutions.

In the past two decades, fractional calculus theory gained a great attention and popularity in various fields of science and engineering due to its demonstrated applications. These contributions to the fields of science and engineering are based on the mathematical analysis. It covers the widely known classical fields such as Abel's integral equation and viscoelasticity. Also, including the analysis of feedback amplifiers, fractional-order Chua-Hartley systems, electrode-electrolyte interface models, fractional-order models of neurons, electric conductance of biological systems, generalized voltage dividers, fitting of experimental data, capacitor theory, and the fields of special functions [14, 17–19, 48, 49].

---

Received: 14 February 2024 ; Accepted: 24 March 2024.

\* Corresponding author. Email: it.kazakov@yandex.ru .

Several robust methods have been used to solve FDEs, fractional integro-differential equations and dynamic systems containing fractional derivatives. Some of the most important methods are the Adomian's decomposition [12, 43, 44], the exp-function [15], the He's the variational iteration [45, 51], the fractional sub-equation [55], the first integral [29], the homotopy analysis [7, 8, 23], the (G'/G)-expansion [3], the generalized tanh-coth [37], the  $\tan(\phi/2)$ -expansion [26], the homotopy perturbation [22, 41], spectral methods [13], the transform [25], and other methods [5, 6, 11, 40, 47, 50].

A substantial amount of research work has been directed for the study of the nonlinear fractional Zakharov-Kuznetsov, KdV and K(2,2) equations given by

$$\frac{\partial^\alpha}{\partial t^\alpha} u + \frac{\partial}{\partial x} u^2 + \frac{1}{8} \frac{\partial^3}{\partial x^3} u^2 + \frac{1}{8} \frac{\partial^3}{\partial x \partial y^2} u^2 = 0, \quad (1.1)$$

and

$$\frac{\partial^\alpha}{\partial t^\alpha} u + \frac{\partial}{\partial x} u^3 + 2 \frac{\partial^3}{\partial x^3} u^3 + 2 \frac{\partial^3}{\partial x \partial y^2} u^3 = 0, \quad (1.2)$$

$$\frac{\partial^\alpha}{\partial t^\alpha} u - 3 \frac{\partial}{\partial x} u^2 + \frac{\partial^3}{\partial x^3} u = 0, \quad (1.3)$$

$$\frac{\partial^\alpha}{\partial t^\alpha} u + \frac{\partial}{\partial x} u^2 + \frac{\partial^3}{\partial x^3} u^2 = 0. \quad (1.4)$$

The ZK equation, first obtained as a description of weakly nonlinear ion-acoustic modes in a strongly magnetized plasma, is of particular interest as it is the simplest equation that admits cylindrical and spherical solitary wave solutions in addition to the planar KdV soliton solutions [52]. Another powerful analytical method is called the Exp-function method (EFM), which was first presented by He [20]. The EFM has successfully been applied to many situations. For example, He [20] solved the nonlinear wave equations via the EFM. Abdou [1] solved generalized solitary and periodic solutions for nonlinear partial differential equations by the EFM. For further information refer to vigorous references therein ([4, 9, 10, 30–33]).

We will use the Jumarie's modified Riemann-Liouville derivative [24] of order  $\alpha$  where is defined by the following expression:

$$D_t^\alpha u(t) = \begin{cases} \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-\tau)^{-\alpha} (u(\tau) - u(0)) d\tau, & \text{if } 0 < \alpha \leq 1, \\ [u^{(n)}(t)]^{(\alpha-n)}, & \text{if } n \leq \alpha < n+1, n \geq 1. \end{cases} \quad (1.5)$$

The properties of the modified Riemann-Liouville derivative are given as:

- (1)  $D^\alpha[f(t)g(t)] = f(t)D^\alpha g(t) + g(t)D^\alpha f(t)$ ,
- (2)  $D^\alpha[f(g(t))] = f'_g(g(t))D^\alpha g(t)$ ,
- (3)  $D^\alpha[f(g(t))] = D_g^\alpha f(g(t))[g'(t)]^\alpha$ ,
- (4)  $D_t^\alpha t^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(1+\alpha-\gamma)} t^{\gamma-\alpha}$ ,  $\gamma > 0$ , where  $\Gamma$  denotes the Gamma function.

Authors of [39] explained the generalized fifth-order KdV like equation with prime number  $p = 3$  via a generalized bilinear differential operator. N-lump was investigated to the variable-coefficient Caudrey-Dodd-Gibbon-Kotera-Sawada equation [38]. Applications of  $\tan(\phi/2)$ -expansion method for the Biswas-Milovic equation [34], the Gerdjikov-Ivanov model [36], the Kundu-Eckhaus equation [35] and the fifth-order integrable equations [27] were studied. Lump solutions were analyzed to the fractional generalized CBS-BK equation [53] and the (3+1)-D Burger system [16]. The approximations of one-dimensional hyperbolic equation with non-local integral conditions were constructed by reduced differential transform method [46]. The generalized Hirota bilinear strategy by the number prime was used to the (2+1)-dimensional generalized fifth-order KdV like equation [39]. The traveling wave solutions and analytical treatment of the simplified MCH equation and the combined KdV-mKdV equations were studied [2].

Our objective here is to find exact solutions of some fractional nonlinear partial differential equations (FNLPEs) under consideration the improved EFM for obtaining the new periodic solitary wave solutions. Discussion about the Improved Exp-function Method is given. Application of the improved EFM on FNLPEs are investigated and derived



exact solutions. In the continuation, we will present graphical illustrations of some solutions of the aforementioned models. After that, we will deal with the investigation of solutions and we will end with a conclusion.

## 2. IMPROVED EXP-FUNCTION METHOD

To illustrate the basic idea of the improved EFM, we take the following nonlinear fractional partial differential equation in the form:

$$\mathcal{N}(u, u_x, u_y, u_{xx}, u_{yy}, \dots, D_t^\alpha u, D_x^\alpha u, D_{xx}^\alpha u, \dots) = 0, \quad 0 < \alpha \leq 1. \tag{2.1}$$

Using a transformation

$$u(x, y, t) = u(\eta), \quad \eta = kx + my + \frac{nt^\alpha}{\Gamma(\alpha + 1)}, \tag{2.2}$$

where  $k, m$  and  $n$  are constants to be determined later, we can rewrite equation Eq. (2.1) in the following nonlinear ODE

$$\mathcal{M}(u, ku', mu', k^2u'', m^2u'', \dots, nu', \dots) = 0, \tag{2.3}$$

where the prime denotes derivative with respect to  $\eta$ . If possible, integrate Eq. (2.3) term by term one or more times. This yields constants of integration. For simplicity, the integration constants can be set to zero. Based on the modified Exp-function method, the final solution can be presented as

$$u(\eta) = \frac{\sum_{n=1}^{2M} a_n \exp(n\eta)}{\sum_{n=1}^{2M} b_n \exp(n\eta)} + \frac{\sum_{n=1}^{2M} a_{-n} \exp(-n\eta)}{\sum_{n=1}^{2M} b_{-n} \exp(-n\eta)}, \tag{2.4}$$

where  $M$  is integers which are unknown to be determine,  $a_n, b_n, a_{-n}$  and  $b_{-n}$  are unfound constants. To determine the value of  $M$ , we balance the linear term of highest order of Eq. (2.3) with the highest order nonlinear term. Plugging (2.4) into the Eq. (2.3), equating the coefficients of each power to zero gives the system of the algebraic equations for  $a_n, b_n, a_{-n}$  and  $b_{-n}$ , then solve the system to determine these constants.

## 3. TEST PROBLEMS

In this section, we offer several examples to demonstrate the applicability of improved EFM to solve FNLPPDEs.

**3.1. The fractional ZK(2,2,2) equation.** Consider the (2+1)-dimensional ZK(2,2,2) equation of fractional order as follows [42]

$$\frac{\partial^\alpha}{\partial t^\alpha} u + \frac{\partial}{\partial x} u^2 + \frac{1}{8} \frac{\partial^3}{\partial x^3} u^2 + \frac{1}{8} \frac{\partial^3}{\partial x \partial y^2} u^2 = 0, \tag{3.1}$$

by utilizing the transformation  $\eta = kx + my + \frac{nt^\alpha}{\Gamma(\alpha+1)}$ , Eq. (3.1) is reduced to an ODE as

$$nu' + k(u^2)' + \frac{1}{8}k^3(u^2)''' + \frac{1}{8}km^2(u^2)''' = 0. \tag{3.2}$$

Integrating Eq. (3.2) once and setting the constant of integration equal to zero, results in

$$nu + ku^2 + \frac{1}{8}(k^3 + km^2)(u^2)'' = 0. \tag{3.3}$$

Balancing the  $(u^2)''$  and  $u$  by employing the homogenous principle, we get

$$2M + 2 = M, \quad \Rightarrow M = -2. \tag{3.4}$$

To get a closed form solution, we use the transformation

$$u(\eta) = v(\eta)^{-2}. \tag{3.5}$$

Substituting (3.5) into Eq. (3.3), we get

$$kv^2 + nv^4 + \frac{1}{2}(k^3 + km^2)(5v'^2 - vv'') = 0. \tag{3.6}$$



Balancing the  $vv''$  and  $v^4$ , we get

$$2M + 2 = 4M, \quad \Rightarrow M = 1. \quad (3.7)$$

Then the exact solution will be as

$$v(\eta) = \frac{a_1 \exp(\eta) + a_2 \exp(2\eta)}{b_1 \exp(\eta) + b_2 \exp(2\eta)} + \frac{a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{b_{-1} \exp(-\eta) + b_{-2} \exp(-2\eta)}. \quad (3.8)$$

For simplicity, we set  $a_2 = b_2 = 1$  and  $a_{-2} = b_{-2} = 1$ , then (3.8) reduces to

$$v(\eta) = \frac{a_1 + \exp(\eta)}{b_1 + \exp(\eta)} + \frac{a_{-1} + \exp(-\eta)}{b_{-1} + \exp(-\eta)}. \quad (3.9)$$

Inserting (3.9) in to Eq. (3.6), we obtain

$$\left( (b_1 + \exp(\eta))^4 (b_{-1} + \exp(-\eta))^4 \right)^{-1} \sum_{n=-4}^4 C_n \exp(n\xi) = 0, \quad (3.10)$$

where  $C_n$  ( $-4 \leq n \leq 4$ ) are polynomial statements in terms of  $a_1, a_{-1}, b_1, b_{-1}, k, n$  and  $m$ . Hence, solving the resulting system  $C_n = 0$  ( $-4 \leq n \leq 4$ ) simultaneously, we acquire the below set of parameters of solutions

**Set I:**

$$a_1 = -b_1, \quad a_{-1} = \frac{1}{b_1}, \quad b_{-1} = -\frac{1}{b_1}, \quad b_1 = b_1, \quad k = \sqrt{\frac{1+2m^2}{2}}i, \quad n = \frac{3}{16}k, \quad m = m. \quad (3.11)$$

We, therefore, gained the following generalized solitary solution

$$u_1(\eta) = \left( \frac{4b_1 e^\eta}{e^{2\eta} - b_1^2} \right)^{-2}, \quad \eta = \sqrt{\frac{1+2m^2}{2}}ix + my + \frac{3}{16} \sqrt{\frac{1+2m^2}{2}}i \frac{t^\alpha}{\Gamma(\alpha+1)}. \quad (3.12)$$

If we choose  $b_1 = 1$ , then Eq. (3.12) get to

$$u_1(x, t) = \frac{1}{4} \sinh^2 \left( \sqrt{\frac{1+2m^2}{2}}ix + my + \frac{3}{16} \sqrt{\frac{1+2m^2}{2}}i \frac{t^\alpha}{\Gamma(\alpha+1)} \right). \quad (3.13)$$

**3.2. The fractional ZK(3,3,3) equation.** As second example, assume the ZK(3,3,3) equation of fractional order [42] be as

$$\frac{\partial^\alpha}{\partial t^\alpha} u + \frac{\partial}{\partial x} u^3 + 2 \frac{\partial^3}{\partial x^3} u^3 + 2 \frac{\partial^3}{\partial x \partial y^2} u^3 = 0, \quad (3.14)$$

by using the transformation  $\eta = kx + my + \frac{nt^\alpha}{\Gamma(\alpha+1)}$ , Eq. (3.14) is reduced to an ODE as

$$nu' + k(u^3)' + 2k^3(u^3)''' + 2km^2(u^3)''' = 0. \quad (3.15)$$

Integrating Eq. (3.15) once and setting the constant of integration equal to zero, results in

$$nu + ku^3 + 2(k^3 + km^2)(u^3)'' = 0. \quad (3.16)$$

Balancing the  $(u^3)''$  and  $u$ , we achieve to  $M = -1$ . To obtain an exact solution, we get

$$u(\eta) = v(\eta)^{-1}. \quad (3.17)$$

Plugging (3.17) into Eq. (3.16), we obtain

$$kv^2 + nv^4 + 6(k^3 + km^2)(4v'^2 - vv'') = 0. \quad (3.18)$$

Balancing the  $vv''$  and  $v^4$ , we get to  $M = 1$ . Therefore, the exact solution will be as

$$v(\eta) = \frac{a_1 \exp(\eta) + a_2 \exp(2\eta)}{b_1 \exp(\eta) + b_2 \exp(2\eta)} + \frac{a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{b_{-1} \exp(-\eta) + b_{-2} \exp(-2\eta)}. \quad (3.19)$$






FIGURE 1. 3D and 2D plots of (3.13) (a)  $\alpha = 0.5$ , (c)  $\alpha = 0.9$  when  $-100 < x < 100$  and  $0 < t < 10$  and (b)  $\alpha = 0.5$ , (d)  $\alpha = 0.9$  when  $y = 1$ ,  $m = 2$ ,  $-100 < x < 100$  and  $t = 10$ .

For simplicity, we set  $a_2 = b_2 = 1$  and  $a_{-2} = b_{-2} = 1$ , then (3.19) transforms to

$$v(\eta) = \frac{a_1 + \exp(\eta)}{b_1 + \exp(\eta)} + \frac{a_{-1} + \exp(-\eta)}{b_{-1} + \exp(-\eta)}. \tag{3.20}$$

Substituting (3.20) into Eq. (3.18), we acquire the below set of solutions

**Set I:**

$$a_1 = -b_1, \quad a_{-1} = \frac{1}{b_1}, \quad b_{-1} = -\frac{1}{b_1}, \quad b_1 = b_1, \quad k = \frac{1}{3}\sqrt{\frac{1+18m^2}{2}}i, \quad n = \frac{1}{6}k, \quad m = m, \tag{3.21}$$

$$u_1(x, t) = \left( \frac{4b_1}{b_1^2 e^{-\frac{1}{3}\sqrt{\frac{1+18m^2}{2}}ix - my - \frac{1}{18}\sqrt{\frac{1+18m^2}{2}}i\frac{t^\alpha}{\Gamma(\alpha+1)}} - e^{\frac{1}{3}\sqrt{\frac{1+18m^2}{2}}ix + my + \frac{1}{18}\sqrt{\frac{1+18m^2}{2}}i\frac{t^\alpha}{\Gamma(\alpha+1)}}} \right)^{-1}. \tag{3.22}$$

If we choose  $b_1 = 1$ , then Equation (3.22) achieve to

$$u_1(x, t) = \frac{1}{4} \sinh \left( \frac{1}{3}\sqrt{\frac{1+18m^2}{2}}ix + my + \frac{1}{18}\sqrt{\frac{1+18m^2}{2}}i\frac{t^\alpha}{\Gamma(\alpha+1)} \right). \tag{3.23}$$






FIGURE 2. 3D and 2D plots of (3.23) (a)  $\alpha = 0.5$ , (c)  $\alpha = 0.9$  when  $-100 < x < 100$  and  $0 < t < 10$  and (b)  $\alpha = 0.5$ , (d)  $\alpha = 0.9$  when  $y = 1$ ,  $m = 2$ ,  $-100 < x < 100$  and  $t = 10$ .

**3.3. The fractional KdV equation.** As third example, we next consider (1+1)-dimensional fractional KdV equation [7]

$$\frac{\partial^\alpha}{\partial t^\alpha} u - 3 \frac{\partial}{\partial x} u^2 + \frac{\partial^3}{\partial x^3} u = 0, \quad (3.24)$$

by using the transformation  $\eta = kx + \frac{nt^\alpha}{\Gamma(\alpha+1)}$ , Eq. (3.24) is reduced to an ODE as

$$nu' - 3k(u^2)' + k^3(u)''' = 0. \quad (3.25)$$

Integrating Eq. (3.25) once and setting the constant of integration equal to zero, concludes

$$nu - 3ku^2 + k^3u'' = 0. \quad (3.26)$$

Balancing the  $u''$  and  $u^2$ , we obtain  $M = 2$ . Therefore, the exact solution can be written as

$$u(\eta) = \frac{\sum_{n=1}^4 a_n \exp(n\eta)}{\sum_{n=1}^4 b_n \exp(n\eta)} + \frac{\sum_{n=1}^4 a_{-n} \exp(-n\eta)}{\sum_{n=1}^4 b_{-n} \exp(-n\eta)}. \quad (3.27)$$

For simplicity, we set  $a_3 = b_3 = a_4 = b_4 = 1$  and  $a_{-3} = b_{-3} = a_{-4} = b_{-4} = 1$ , then (3.27) reduces to

$$u(\eta) = \frac{a_1 + a_2 e^\eta + e^{2\eta} + e^{3\eta}}{b_1 + b_2 e^\eta + e^{2\eta} + e^{3\eta}} + \frac{a_{-1} + a_{-2} e^{-\eta} + e^{-2\eta} + e^{-3\eta}}{b_{-1} + b_{-2} e^{-\eta} + e^{-2\eta} + e^{-3\eta}}. \quad (3.28)$$



Plugging (3.28) into Eq. (3.26), we gain the below set of solutions as

**Set I:**

$$a_{-1} = a_{-2} = \frac{4k^2 - 3}{3b_1}, \quad a_1 = a_2 = \frac{1}{3}(4k^2 - 3)b_1, \quad b_{-1} = b_{-2} = \frac{1}{b_1}, \quad b_1 = b_1, \quad k = k, \quad n = 4k^3, \quad (3.29)$$

$$u_1(x, t) = \frac{4}{3} \frac{k^2 \left[ b_1^2 e^{-2k\left(x + \frac{4k^2 t^\alpha}{\Gamma(\alpha+1)}\right)} + e^{2k\left(x + \frac{4k^2 t^\alpha}{\Gamma(\alpha+1)}\right)} + 2b_1 \right]}{\left[ e^{-2k\left(x + \frac{4k^2 t^\alpha}{\Gamma(\alpha+1)}\right)} b_1 + 1 \right] \left[ e^{2k\left(x + \frac{4k^2 t^\alpha}{\Gamma(\alpha+1)}\right)} + b_1 \right]}. \quad (3.30)$$

**Set II:**

$$a_1 = a_2 = \frac{a_{-1}}{b_{-1}^2}, \quad a_{-1} = a_{-1} = a_{-2}, \quad b_{-1} = b_{-1} = b_{-2}, \quad b_1 = b_2 = \frac{1}{b_{-1}}, \quad k = k, \quad n = \frac{3k(b_{-1} + a_{-1})}{b_{-1}}, \quad (3.31)$$

$$u_2(x, t) = \frac{1}{b_{-1}} \left[ \frac{(a_{-1} + b_{-1})e^{-\lambda} + 2b_{-1}^2 + a_{-1}b_{-1}^2 e^\lambda + b_{-1}^3 e^\lambda}{(e^{-\lambda} + b_{-1})(b_{-1} e^\lambda + 1)} \right], \quad (3.32)$$

where  $\lambda = \frac{2k}{b_{-1}} \left( x b_{-1} + 3(b_{-1} + a_{-1}) \frac{t^\alpha}{\Gamma(\alpha+1)} \right)$ . If we choose  $b_{-1} = a_{-1} = 1$ , then Equation (3.32) get to

$$u_2(x, t) = \frac{e^{-\lambda} + 2 + 2e^\lambda}{e^{-\lambda} + 2 + e^\lambda}, \quad \lambda = 2kx + \frac{12kt^\alpha}{\Gamma(\alpha + 1)}. \quad (3.33)$$

**Set III:**

$$a_{-1} = a_{-2} = \frac{3k^2 - 3}{b_1}, \quad a_1 = a_2 = (3k^2 - 1)b_1, \quad b_{-1} = b_{-2} = \frac{1}{b_1}, \quad b_1 = b_1, \quad k = k, \quad n = 9k^3, \quad (3.34)$$

$$u_3(x, t) = \frac{3k^2 \left[ b_1^3 e^{-2k\left(x + 9k^2 \frac{t^\alpha}{\Gamma(\alpha+1)}\right)} + e^{2k\left(x + 9k^2 \frac{t^\alpha}{\Gamma(\alpha+1)}\right)} + 2b_1 \right]}{\left[ e^{-2k\left(x + 9k^2 \frac{t^\alpha}{\Gamma(\alpha+1)}\right)} b_1 + 1 \right] \left[ e^{2k\left(x + 9k^2 \frac{t^\alpha}{\Gamma(\alpha+1)}\right)} + b_1 \right]}. \quad (3.35)$$

**3.4. The fractional K(2,2) equation.** The (1+1)-dimensional K(2,2) equation of fractional order is given as

$$\frac{\partial^\alpha}{\partial t^\alpha} u + \frac{\partial}{\partial x} u^2 + \frac{\partial^3}{\partial x^3} u^2 = 0, \quad (3.36)$$

by using the transformation  $\eta = kx + \frac{nt^\alpha}{\Gamma(\alpha+1)}$ , Eq. (3.36) is reduced to an ODE as

$$nu' + k(u^2)' + k^3(u^2)''' = 0. \quad (3.37)$$

Integrating Eq. (3.37) once and setting the constant of integration equal to zero, concludes

$$nu + ku^2 + k^3(u^2)'' = 0. \quad (3.38)$$

Balancing the  $(u^2)''$  and  $u$ , we get  $M = -2$ . To get an exact solution, we obtain

$$u(\eta) = v(\eta)^{-2}. \quad (3.39)$$

Substituting Eq. (3.39) into Eq. (3.38), we get

$$kv^2 + nv^4 + 4k^3(5v'^2 - vv'') = 0. \quad (3.40)$$

Balancing the  $vv''$  and  $v^4$ , we achieve to  $M = 1$ . Then the exact solution is as

$$v(\eta) = \frac{a_1 \exp(\eta) + a_2 \exp(2\eta)}{b_1 \exp(\eta) + b_2 \exp(2\eta)} + \frac{a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{b_{-1} \exp(-\eta) + b_{-2} \exp(-2\eta)}. \quad (3.41)$$






FIGURE 3. 3D and 2D plots of (3.33) (a)  $\alpha = 0.5$ , (b)  $\alpha = 0.9$  when  $0 < x < 10$  and  $-100 < t < 100$  and (c)  $\alpha = 0.5$ , (d)  $\alpha = 0.9$  when  $k = 2$ ,  $-100 < t < 100$  and  $x = 0.5$ .

For simplicity, we set  $a_2 = b_2 = 1$  and  $a_{-2} = b_{-2} = 1$ , then Eq. (3.41) reduces to

$$v(\eta) = \frac{a_1 \exp(\eta) + \exp(2\eta)}{b_1 \exp(\eta) + \exp(2\eta)} + \frac{a_{-1} \exp(-\eta) + \exp(-2\eta)}{b_{-1} \exp(-\eta) + \exp(-2\eta)}. \quad (3.42)$$

Plugging (3.42) into Eq. (3.40), we gain the following set of solutions as

**Set I:**

$$a_1 = -b_1, \quad a_{-1} = \frac{1}{b_1}, \quad b_{-1} = -\frac{1}{b_1}, \quad b_1 = b_1, \quad k = \frac{1}{4}i, \quad n = \frac{3}{64}i, \quad (3.43)$$

$$u_1(x, t) = \frac{1}{16b_1^2} \left( b_1 + e^{\frac{i}{64}[16x+3\frac{t^\alpha}{\Gamma(\alpha+1)}]} \right)^2 \left( -1 + b_1 e^{-\frac{i}{64}[16x+3\frac{t^\alpha}{\Gamma(\alpha+1)}]} \right)^2. \quad (3.44)$$

If we choose  $b_1 = 1$ , then Equation (3.44) get to

$$u_1(x, t) = -\frac{I}{8} \sin \left[ \frac{1}{64} \left( 16x + \frac{3t^\alpha}{\Gamma(\alpha+1)} \right) \right], \quad I = \sqrt{-1}. \quad (3.45)$$

#### 4. PHYSICAL INTERPRETATIONS OF THE SOLUTIONS

Numerical simulations have been performed by using the Maple software. We depict some of the solutions to have an idea on the mechanism of the original equations. Particularly, we depict solutions of the fractional ZK(2,2,2), ZK(3,3,3), KdV and K(2,2) equations by considering the suitable values of the parameters obtained. The graphical






FIGURE 4. 3D and 2D plots of (3.45) (a)  $\alpha = 0.5$ , (b)  $\alpha = 0.9$  when  $0 < x < 10$  and  $-100 < t < 100$  and (c)  $\alpha = 0.5$ , (d)  $\alpha = 0.9$  when  $-100 < t < 100$  and  $x = 0.5$ .

representations to this solution are presented in Figures 1-4, respectively. Figures 1-4 represent the variation of some appropriated parameters. The curves 1-4 have been plotted for different values of appropriated parameters. In Figures 1-4, we plot two and three dimensional graphics of real values of (3.13), (3.23), (3.33), and (3.45), respectively.

## 5. CONCLUSION

In this paper, we employed the modified Exp-function method for deriving the exact solitary wave solutions for some of the fractional partial differential equations. As a consequence, we gained many new exact solitary solutions for the aforementioned equations which are expressed by a rational exponential function form. The modified Exp-function method has many applications in the field of engineering, mathematical science, and physics for solving a large class of non-linear partial differential equations. We then think that these results will help for conducting future research in diverse areas of physics such as mathematical physics, nonlinear phenomena, fluid mechanics and other applied fields and so on. The method utilized here can be applied to other nonlinear fractional partial differential equations.

## REFERENCES

- [1] M. A. Abdou, *Generalized solitary and periodic solutions for nonlinear partial differential equations by the Exp-function method*, *Nonlinear Dyn.*, 512 (2008), 1-9.
- [2] N. H. Ali, S. A. Mohammed, and J. Manafian, *Study on the simplified MCH equation and the combined KdV-mKdV equations with solitary wave solutions*, *Partial Diff. Eq. Appl. Math.*, 9 (2024), 100599.
- [3] A. Bekir, O. Guner, A. H. Bhrawy, and A. Biswas, *Solving nonlinear fractional differential equations using exp-function and  $(G'/G)$ -expansion methods*, *Rom. J. Phys.*, 60(3-4) (2015), 360-378.



- [4] A. Boz and A. Bekir, *Application of Exp-function method for (3+1)-dimensional nonlinear evolution equations*, *Comput. Math. Appl.*, *56* (2008), 1451-1456.
- [5] R. Churchill, *Operational Mathematics (3rd edition)*, McGraw-Hill, New York, 1972.
- [6] L. Debanth, *Recent applications of fractional calculus to science and engineering*, *Int. J. Math. Math. Sci.*, *54* (2003), 3413-3442.
- [7] M. Dehghan, J. Manafian, and A. Saadatmandi, *Solving nonlinear fractional partial differential equations using the homotopy analysis method*, *Num. Meth. Partial Differential Eq J.*, *26* (2010), 448-479.
- [8] M. Dehghan, J. Manafian, and A. Saadatmandi, *The solution of the linear fractional partial differential equations using the homotopy analysis method*, *Z. Naturforsch.*, *65(a)* (2010), 935-949.
- [9] M. Dehghan, J. Manafian Heris, and A. Saadatmandi, *Analytical treatment of some partial differential equations arising in mathematical physics by using the Exp-function method*, *Int. J. Modern Phys. B*, *25* (2011), 2965-2981.
- [10] M. Dehghan, J. Manafian Heris, and A. Saadatmandi, *Application of the Exp-function method for solving a partial differential equation arising in biology and population genetics*, *Int. J. Num. Methods for Heat Fluid Flow*, *21* (2011), 736-753.
- [11] D. G. Duffy, *Transform Methods for Solving Partial Differential*, Chapman and Hall/CRC, 2004.
- [12] V. D. Gejji and H. Jafari, *Solving a multi-order fractional differential equation*, *Appl. Math. Comput*, *189* (2007), 541-548.
- [13] K. A. Gepreel and M. S. Mohamed, *An optimal homotopy analysis method nonlinear fractional differential equation*, *Journal of Advanced Research in Dynamical and Control Systems*, *6(1)* (2014), 1-10.
- [14] I. Grigorenko and E. Grigorenko, *Chaotic dynamics of the fractional lorenz system*, *Phys. Rev. Lett*, *91* (2003), 034101. 2007) 541-548.
- [15] O. Guner, A. Bekir, and H. Bilgil, *A note on exp-function method combined with complex transform method applied to fractional differential equations*, *Adv. Nonlinear Anal.*, *4(3)* (2015), 201-208.
- [16] Y. Gu, S. Malmir, J. Manafian, O. A. Ilhan, A. A. Alizadeh, and A. J. Othman, *Variety interaction between k-lump and k-kink solutions for the (3+1)-D Burger system by bilinear analysis*, *Results Phys.*, *43* (2022), 106032.
- [17] J. He, *Nonlinear oscillation with fractional derivative and its applications*, *International Conference on Vibrating Engineering*, Dalian, China, (1998), 288-291.
- [18] J. He, *Approximate analytical solution for seepage flow with fractional derivatives in porous media*, *Comput. Methods. Appl. Mech. Eng.*, *167* (1998), 57-68.
- [19] J. He, *Some applications of nonlinear fractional differential equations and their approximations*, *Bull. Sci. Technol.*, *15* (1999), 86-90.
- [20] J. H. He and X. H. Wu, *Exp-function method for nonlinear wave equations*, *Chaos, Solitons Fractals*, *30* (2006), 700-708.
- [21] J. H. He, S. K. Elagan, and Z. B. Li, *Geometrical explanation of the fractional complex transform and derivative chain rule for fractional calculus*, *Phys. Lett. A*, *376* (2012), 257-259.
- [22] M. A. Herzallah and K. A. Gepreel, *Approximate solution to time-space fractional cubic nonlinear Schrodinger equation*, *Applied Mathematical Modeling*, *36(11)* (2012), 5678-5685.
- [23] H. Jafari and S. Seifi, *Solving a system of nonlinear fractional partial differential equations using homotopy analysis method*, *Commu. Nonlinear Sci. Num. Simu*, *14* (2009), 1962-1969.
- [24] G. Jumarie, *Modified Riemann-Liouville derivative and fractional Taylor series of nondifferentiable functions further results*, *Comput. Math. Appl.*, *51* (2006), 1367-1376.
- [25] S. Kemple and H. Beyer, *Global and causal solutions of fractional differential equations*, *Transform methods and special functions: Varna 96*, *Proceedings of 2nd international workshop (SCTP)*, Singapore, *96* (1997), 210-216.
- [26] M. Lakestani and J. Manafian, *Analytical treatment of nonlinear conformable timefractional Boussinesq equations by three integration methods*, *Opt Quant Electron*, *50(4)* (2018), 1-31.



- [27] M. Lakestani, J. Manafian, A. R. Najafizadeh, and M. Partohaghghi, *Some new soliton solutions for the nonlinear the fifth-order integrable equations*, *Comput. Meth. Diff. Equ.*, *10*(2) (2022), 445-460.
- [28] Z. B. Li and J. H. He, *Application of the fractional complex transform to fractional differential equations*, *Nonlinear Sci. Lett. A*, *2* (2011), 121-126.
- [29] B. Lu, *The first integral method for some time fractional differential equations*, *J. Math. Anal. Appl.*, *395* (2012), 684-693.
- [30] J. Manafian Heris and M. Bagheri, *Exact solutions for the modified KdV and the generalized KdV equations via Exp-function method*, *J. Math. Extension*, *4* (2010), 77-98.
- [31] J. Manafian Heris and M. Fazli Aghdaei, *Application of the Exp-function method for solving the combined KdV-mKdV and Gardner-KP equations*, *Math. Science*, *6* (2012), 68.
- [32] J. Manafian Heris and I. Zamanpour, *Analytical treatment of the coupled Higgs equation and the Maccari system via Exp-Function Method*, *Acta Universitatis Apulensis*, *33* (2013), 203-216.
- [33] J. Manafian Heris and M. Lakestani, *Solitary wave and periodic wave solutions for variants of the KdV-Burger and the  $K(n,n)$ -Burger equations by the generalized tanh-coth method*, *Commu. Num. Anal.*, *2013* (2013), 1-18.
- [34] J. Manafian and M. Lakestani, *Application of  $\tan(\phi/2)$ -expansion method for solving the Biswas-Milovic equation for Kerr law nonlinearity*, *Optik*, *127*(4) (2016), 2040-2054.
- [35] J. Manafian and M. Lakestani, *Abundant soliton solutions for the Kundu-Eckhaus equation via  $\tan(\phi(\xi))$ -expansion method*, *Optik*, *127*(14) (2016), 5543-5551.
- [36] J. Manafian and M. Lakestani, *Optical soliton solutions for the Gerdjikov-Ivanov model via  $\tan(\phi/2)$ -expansion method*, *Optik*, *127*(20) (2016), 9603-9620.
- [37] J. Manafian and M. Lakestani, *A new analytical approach to solve some of the fractional-order partial differential equations*, *Indian J Phys.*, *91*(3) (2016), 243-258.
- [38] J. Manafian and M. Lakestani,  *$N$ -lump and interaction solutions of localized waves to the  $(2+1)$ - dimensional variable-coefficient Caudrey-Dodd-Gibbon-Kotera-Sawada equation*, *J. Geom. Phys.*, *150* (2020), 103598.
- [39] J. Manafian, L. A. Dawood, and M. Lakestani, *New solutions to a generalized fifth-order KdV like equation with prime number  $p = 3$  via a generalized bilinear differential operator*, *Partial Diff. Eq. Appl. Math.*, *9* (2024), 100600.
- [40] K. S. Miller and B. Ross, *An introduction to the fractional calculus and fractional differential equations*, New York, Wiley, 1993.
- [41] M. S. Mohamed, *Analytical treatment of Abel integral equations by optimal homotopy analysis transform method*, *Journal of Information and Computing Science*, *10*(1) (2015), 19-28.
- [42] R. Y. Molliq and B. Batiha, *Approximate analytic solutions of fractional Zakharov-Kuznetsov equations by fractional complex transform*, *Int. J. Eng. Tech.*, *1* (2012), 1-13.
- [43] S. Momani and M. A. Noor, *Numerical methods for fourth-order fractional integrodifferential equations*, *Appl. Math. Comput.*, *182* (2006), 754-760.
- [44] S. Momani and N. T. Shawagfeh, *Decomposition method for solving fractional Riccati differential equations*, *Appl. Math. Comput.*, *182* (2006), 1083-1092.
- [45] S. Momani and Z. Odibat, *Homotopy perturbation method for nonlinear partial differential equations of fractional order*, *Phys. Lett. A*, *365* (2007), 345-350.
- [46] S. R. Moosavi, N. Taghizadeh, and J. Manafian, *Analytical approximations of one-dimensional hyperbolic equation with non-local integral conditions by reduced differential transform method*, *Comput. Meth. Diff. Equ.*, *8*(3) (2020), 537-552.
- [47] I. Podlubny, *Fractional differential equations: An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*, New York: Academic Press, 1999.
- [48] I. Podlubny, *Fractional differential equations*, Academic Press, San Diego, 1999.
- [49] I. Podlubny, *Geometric and physical interpretation of fractional integration and fractional differentiation*, *Fract. Calculus. Appl. Anal.*, *5* (2002), 367-386.



- [50] S. G. Samko, A. Kilbas, and O. Marichev, *Fractional integrals and derivatives: theory and applications*, Yverdon, Gordon and Breach, 1993.
- [51] N. H. Sweilam, M. M. Khader, and R. F. Al-Bar, *Numerical studies for a multi-order fractional differential equation*, Phys. Lett. A, *371* (2007), 26-33.
- [52] V. E. Zakharov and E. A. Kuznetsov, *On three-dimensional solitons*, Sov. Phys, *39* (1974), 285-288.
- [53] M. Zhang, X. Xie, J. Manafian, O. A. Ilhan, and G. Singh, *Characteristics of the new multiple rogue wave solutions to the fractional generalized CBS-BK equation*, J. Adv. Res., *38* (2022), 131-142.
- [54] X. Zhao, H. Zhou, Y. Tang, and H. Jia, *Travelling wave solutions for modified Zakharov-Kuznetsov equation*, Appl. Math. Comput, *181* (2006), 634-648.
- [55] B. Zheng and C. Wen, *Exact solutions for fractional partial differential equations by a new fractional sub-equation method*, Advances in Difference Equations, *2013* (2013), 199.

