



Dynamical systems in product Łukasiewicz semirings

Ali Molkhasi^{1,*} and Mahsa Ezzati²

Department of Mathematics, Faculty of Mathematical Sciences, Farhangian University, Tehran, Iran.

Abstract

This paper studies about dynamical systems in product Łukasiewicz semirings and we generalize the results of Markechova and Riečan concerning the logical entropy. Also, the notion of logical entropy of a product Łukasiewicz semiring is introduced and it is shown that entropy measure is invariant under isomorphism.

Keywords. Łukasiewicz semiring, Entropy, Dynamical systems, Partition.

2010 Mathematics Subject Classification. 65L05, 34K06, 34K28.

1. INTRODUCTION

Shannon entropy in [25] concept of information theory is studied. Kolmogorov and Sinai [11, 26] developed the entropy of a dynamical system. Recall that if two dynamical systems are isomorphic, then they have the same entropy. Recently in paper [17], in a dynamical system, the notion of logical entropy is studied and it is shown that by replacing the Shannon entropy function by the logical entropy function are obtained the results that are analogous to the case of classical Kolmogorov-Sinai entropy theory [17, 18]. Here, in a product Łukasiewicz semiring, we introduced logical entropy.

Dynamical systems theory has its origins in the seminal work of Henri Poincar [20] on celestial mechanics. Eventually, dynamical systems theory (or nonlinear dynamics) found broad application beyond celestial mechanics (see, [7]-[16], [24], and [27]). We recall from [19] that there are about entropy of dynamical systems in MV-algebras, especially after its Mundicĭi characterization as an interval in a lattice ordered group. Petrovicovĭ in [21, 22], for the product MV-algebras, a entropy theory of Shannon and Kolmogorov-Sinai type has been provided. Recall that the notion of near semirings, theory of quantum mechanics, and Łukasiewicz near semirings introduced in [2] and [5, 6].

In the present paper, we introduced the notion of logical entropy of a product Łukasiewicz semiring dynamical system and prove entropy measure is invariant under isomorphism of product Łukasiewicz semiring dynamical systems.

2. MAIN RESULTS

Recall that a groupoid is a set having one binary operation satisfying only closure and monoid is a set that is closed under an associative binary operation and has an identity element. We have the following definition from [3]. A near semiring is an algebra $(R, +, \cdot, 0, 1)$ of type $(2, 2, 0, 0)$ such that:

1. $(R, +, 0)$ is a commutative monoid;
2. $(R, \cdot, 1)$ is a groupoid with condition $x \cdot 1 = x = 1 \cdot x$;
3. $(x + y) \cdot z = (x \cdot z) + (y \cdot z)$;
4. $x \cdot 0 = 0 \cdot x = 0$;

Received: 31 July 2023 ; Accepted: 14 May 2024.

* Corresponding author. Email: molkhasi@cfu.ac.ir.

for all $x, y, z \in R$.

Also, a near semiring, which is a monoid, is said a semiring if $(R, \cdot, 1)$ satisfies left distributivity: $x \cdot (y + z) = (x \cdot y) + (x \cdot z)$, for all $x, y, z \in R$. A near semiring R if satisfies in the property $a + a = a$ is called idempotent, for all $a \in R$. It is clear that $(R, +)$ is a semilattice ([3], Remark 1).

By [3], a function α of an idempotent near semiring $(R, +, \cdot, \alpha, 0, 1)$ to $(R, +, \cdot, \alpha, 0, 1)$ is said an involution on R if, for each $a, b \in R$:

1. $a^{\alpha\alpha} = a$;
 2. if $a \leq b$ then $b^\alpha \leq a^\alpha$,
- with \leq the induced order.

Definition 2.1. A product Lukasiewicz semiring is an algebraic structure $(R, +, \cdot, \alpha, \times, 0, 1)$, where $(R, +, \cdot, \alpha, 0, 1)$ is a Lukasiewicz semiring and \times is an associative and abelian binary operation on R with the following properties:

1. For every $x \in R$, $u \times x = x$;
2. If $((x^{\alpha\alpha} + y^\alpha) \cdot y^{\alpha\alpha})^\alpha = 0$, then $z \times x + z \times y \leq u$, and $z \times (x + y) = z \times x + z \times y$, for all $x, y, z \in R$.

We recall that n -tuple $X = (x_1, x_2, \dots, x_n)$ of elements of a product Lukasiewicz semiring $(R, +, \cdot, \alpha, \times, 0, 1)$ with the property $x_1 + x_2 + \dots + x_n = 1$ is called a partition of unity 1. There is the following definition of [21]:

Definition 2.2. A dynamical system in a product Lukasiewicz semiring $(R, +, \cdot, \alpha, \times, 0, 1)$ is a system (R, μ, \mathfrak{U}) , where $\mu : R \rightarrow [0, 1]$ is a state, and $\mathfrak{U} : R \rightarrow R$ is a function such that $\mathfrak{U}(u) = u$, the following conditions are satisfied:

- (i) if $((x^{\alpha\alpha} + y^\alpha) \cdot y^{\alpha\alpha})^\alpha = 0$, then $((\mathfrak{U}(x))^{\alpha\alpha} + (\mathfrak{U}(y))^\alpha) \cdot (\mathfrak{U}(y))^{\alpha\alpha})^\alpha = 0$ and $\mathfrak{U}(x + y) = \mathfrak{U}(x) + \mathfrak{U}(y)$;
 - (ii) $\mathfrak{U}(x \times y) = \mathfrak{U}(x) \times \mathfrak{U}(y)$;
 - (iii) $\mu(\mathfrak{U}(x)) = \mu(x)$;
- for every $x, y \in R$.

Definition 2.3. If $X = (x_1, x_2, \dots, x_n)$ is a partition of unity, then its entropy is defined by the formula

$$H(X) = \sum_{i=1}^n \varphi(\mu(x_i)),$$

where $\varphi(x) = -x \log x$, if $x > 0$, $\varphi(0) = 0$.

Definition 2.4. Suppose (R, μ, \mathfrak{U}) is a dynamical system in a product Lukasiewicz semiring $(R, +, \cdot, \alpha, \times, 0, 1)$ and θ is a sub-additive generator, where a function $\theta : [0, 1] \rightarrow [0, \infty)$ is called a sub-additive generator, if the following implication holds $c_{ij} \in [0, 1]$, $i = 1, \dots, n$, $j = 1, \dots, m$, $\sum_{i=1}^n c_{ij} = b_j$, $\sum_{j=1}^m c_{ij} = a_i$, $\sum_{i=1}^n a_i = 1$, $\sum_{j=1}^m b_j = 1$, then

$$\sum_{i=1}^n \sum_{j=1}^m \theta(c_{ij}) \leq \sum_{i=1}^n \theta(a_i) + \sum_{j=1}^m \theta(b_j).$$

We can define the θ -entropy of (R, μ, \mathfrak{U}) by the formula

$$H_\theta^\mu(\mathfrak{U}) = \sup\{H_\theta^\mu(\mathfrak{U}, X); X \text{ is a partition in } (R, +, \cdot, \alpha, \times, 0, 1)\},$$

where

$$H_\theta^\mu(\mathfrak{U}, X) = \lim_{n \rightarrow \infty} 1/n H_\theta^\mu\left(\bigvee_{k=0}^{n-1} \mathfrak{U}^k(X)\right)$$

Definition 2.5. Suppose $X = (x_1, x_2, \dots, x_n)$ is a partition in a product Lukasiewicz semiring $(R, +, \cdot, \alpha, \times, 0, 1)$, and $\mu : R \rightarrow [0, 1]$ is a state. By Shannon's formula, we define the entropy of X with respect to μ :

$$H_s^\mu(X) = \sum_{i=1}^n s(\mu(x_i)),$$



where $s : [0, 1] \rightarrow [0, \infty)$ is the Shannon entropy function defined by $s(x) = x \log x$ and the Shannon entropy is a number, for every $x \in [0, 1]$. Let $X = (x_1, x_2, \dots, x_n)$ and $Y = (y_1, y_2, \dots, y_n)$ be two partitions in R . Then the conditional entropy of X given Y is defined by

$$H_s^\mu(X/Y) = \sum_{i=1}^n \sum_{j=1}^m (\mu(x_i \times y_j) \cdot \log \frac{\mu(x_i \times y_j)}{\mu(y_j)}).$$

Recall that from [23] the logical entropy of a dynamical system (R, μ, \mathfrak{U}) in a product Lukasiewicz semiring $(R, +, \cdot, \alpha, \times, 0, 1)$ is defined by

$$H_s^\mu(\mathfrak{U}) = \sup\{H_s^\mu(\mathfrak{U}, X); X \text{ is a partition in } R\},$$

and if X is a partition in R , then the logical entropy of \mathfrak{U} relative to X is defined by

$$H_s^\mu(\mathfrak{U}, X) = \lim_{n \rightarrow \infty} H_s^\mu\left(\bigvee_{k=0}^{n-1} \mathfrak{U}^k(X)\right).$$

Theorem 2.6. [18] Let (R, μ, \mathfrak{U}) be a dynamical system in a product Lukasiewicz near semiring $(R, +, \cdot, \alpha, \times, 0, 1)$, and X be a partition in R . For any non-negative integer r , it holds:

$$H_s^\mu(\mathfrak{U}, X) = H_s^\mu\left(\mathfrak{U}, \bigvee_{k=0}^{n-1} \mathfrak{U}^k(X)\right).$$

Definition 2.7. Two product Lukasiewicz near semiring dynamical systems $(R_1, \mu_1, \mathfrak{U}_1)$, $(R_2, \mu_2, \mathfrak{U}_2)$ are isomorphic, if there exists a mapping $\lambda : R_1 \rightarrow R_2$ with $\lambda(u_1) = u_2$ and satisfying the following conditions:

1. $\lambda(x \times y) = \lambda(x) \times \lambda(y)$;
2. If $((x^{\alpha\alpha} + y^\alpha) \cdot y^{\alpha\alpha})^\alpha = 0$, then $\lambda(x + y) = \lambda(x) + \lambda(y)$;
3. $\mu_2(\lambda(x)) = \mu_1(x)$;
4. $\lambda(\mathfrak{U}_1(x)) = \mathfrak{U}_2(\lambda(x))$;

for every $x, y \in \mathfrak{U}_1$.

Now, we are ready to state the following theorem:

Theorem 2.8. Let θ be a sub-additive generator, and $(R_1, \mu_1, \mathfrak{U}_1)$, $(R_2, \mu_2, \mathfrak{U}_2)$ be product Lukasiewicz near semiring dynamical systems. If $\mathfrak{U}_1 \simeq \mathfrak{U}_2$, then $H_\theta^{\mu_1}(\mathfrak{U}_1) = H_\theta^{\mu_2}(\mathfrak{U}_2)$.

Proof. Suppose $\mathfrak{U}_1 \simeq \mathfrak{U}_2$ and λ be an isomorphism between dynamical systems $(R_1, \mu_1, \mathfrak{U}_1)$ and $(R_2, \mu_2, \mathfrak{U}_2)$. Now, assume that $X = (x_1, x_2, \dots, x_n)$ is a partition in a product Lukasiewicz near semiring (R_1, \times) with $x_1 + x_2 + \dots + x_n = u_1$. By definition 2.6, we have $\lambda(x_1 + x_2 + \dots + x_n) = \lambda(x_1) + \lambda(x_2) + \dots + \lambda(x_n) = \lambda(u_1) = u_2$. Therefore, it is show that $\lambda(X) = (\lambda(x_1), \lambda(x_2), \dots, \lambda(x_n))$ in product Lukasiewicz near semiring dynamical system $(R_2, \mu_2, \mathfrak{U}_2)$ is a partition. Also, by definition 2.7,

$$H_\theta^{\mu_2}(\lambda(X)) = \sum_{i=1}^n \theta(\mu_2(\lambda(x_i))) = \sum_{i=1}^n \theta(\mu_1(x_i)) = H_\theta^{\mu_1}(X).$$

On the other hand we have

$$\begin{aligned} H_\theta^{\mu_2}\left(\bigvee_{k=0}^{n-1} \mathfrak{U}_2^k(\lambda(X))\right) &= H_\theta^{\mu_2}\left(\bigvee_{k=0}^{n-1} \lambda(\mathfrak{U}_1^k(X))\right) \\ &= H_\theta^{\mu_2}\left(\lambda\left(\bigvee_{k=0}^{n-1} \mathfrak{U}_1^k(X)\right)\right) = H_\theta^{\mu_1}\left(\bigvee_{k=0}^{n-1} \mathfrak{U}_1^k(X)\right). \end{aligned}$$

If $n \rightarrow \infty$, then we conclude

$$H_\theta^{\mu_2}(\mathfrak{U}_2, \lambda(X)) = \lim_{n \rightarrow \infty} \frac{1}{n} H_\theta^{\mu_2}\left(\bigvee_{k=0}^{n-1} \mathfrak{U}_2^k(\lambda(X))\right)$$



$$= \lim_{n \rightarrow \infty} \frac{1}{n} H_\theta^{\mu_1} \left(\bigvee_{k=0}^{n-1} \mathfrak{U}_1^k(X) \right) = H_\theta^{\mu_1}(\mathfrak{U}_1, X).$$

In fact,

$$H_\theta^{\mu_1}(\mathfrak{U}_1) = \sup\{H_\theta^{\mu_1}(\mathfrak{U}_1, X); X \text{ is a partion in } (R_1, \mu_1, \mathfrak{U}_1)\}$$

$$\leq \sup\{H_\theta^{\mu_2}(\mathfrak{U}_2, Y); Y \text{ is a partion in } (R_2, \mu_2, \mathfrak{U}_2)\} = H_\theta^{\mu_2}(\mathfrak{U}_2).$$

Now, we prove that $H_\theta^{\mu_2}(\mathfrak{U}_2) \leq H_\theta^{\mu_1}(\mathfrak{U}_1)$. To do this, it is enough to define

$$\lambda^{-1} : R_2 \longrightarrow R_1$$

that

$$\lambda^{-1}(x \times y) = \lambda^{-1}(\lambda(a) \times \lambda(b)) = \lambda^{-1}(\lambda(a \times b)) = a \times b = \lambda^{-1}(x) \times \lambda^{-1}(y),$$

where for every $x, y \in R_2$ there exist $a, b \in R_1$ such that $\lambda^{-1}(x) = a$ and $\lambda^{-1}(y) = b$. Also,

$$\lambda^{-1}(x + y) = \lambda^{-1}(\lambda(a) + \lambda(b)) = \lambda^{-1}(\lambda(a + b)) = a + b = \lambda^{-1}(x) + \lambda^{-1}(y),$$

which $x + y \leq u_2$ and $a + b \leq u_1$. In addition, we have

$$\lambda^{-1}(\mathfrak{U}_2(x)) = \lambda^{-1}(\mathfrak{U}_2(\lambda(a))) = \lambda^{-1}(\lambda(\mathfrak{U}_1(a))) = \mathfrak{U}_1(a) = \mathfrak{U}_1(\lambda^{-1}(x)),$$

where $x \in R_2$. Therefore, with the same argument for $H_\theta^{\mu_1}(\mathfrak{U}_1) \leq H_\theta^{\mu_2}(\mathfrak{U}_2)$, we obtain

$$H_\theta^{\mu_2}(\mathfrak{U}_2) \leq H_\theta^{\mu_1}(\mathfrak{U}_1).$$

□

Entropy of a Lukasiewicz near semiring dynamical system (R, μ, \mathfrak{U}) is defined by the

$$\varphi(\mathfrak{U}) = \sup\{\varphi(\mathfrak{U}, X) \mid X \text{ is a partition of unity}\},$$

where

$$\varphi(\mathfrak{U}, X) = \lim_{n \rightarrow \infty} H_n(\mathfrak{U}, X).$$

Definition 2.9. Two Lukasiewicz semiring dynamical systems $(R_1, \mu_1, \mathfrak{U}_1)$, $(R_2, \mu_2, \mathfrak{U}_2)$ are equivalent, if there exists a function $\theta : R_1 \longrightarrow R_2$ satisfying the following conditions:

- (i) θ is a bijection;
- (ii) if $a, b, c \in R_1$, $a = b+c$, then $\theta(a) = \theta(b)+\theta(c)$,
- (iii) $\theta(u_1) = u_2$;
- (iv) $\mu_2(\theta(a)) = \mu_1(a)$ for any $a \in R_1$;
- (v) $\mathfrak{U}_2(\theta(a)) = \theta(\mathfrak{U}_1(a))$ for any $a \in R_1$.

Corollary 2.10. If $(R_1, \mu_1, \mathfrak{U}_1)$, $(R_2, \mu_2, \mathfrak{U}_2)$ are equivalent, then $\varphi(\mathfrak{U}_1) = \varphi(\mathfrak{U}_2)$.

Proof. We know that if X is any partition of u_1 , then $\varphi(x)$ is a partition of u_2 , and $H(X) = H(\varphi(X))$. Suppose that δ is an arbitrary positive number and we get a common refinement \mathcal{C} of $X, \dots, \mathfrak{U}_1^{n-1}(X)$ such that $H_n + \delta > H(\mathcal{C})$. Clearly that $H(\mathcal{C}) \geq H_n(\theta(X))$. Since $H_n(X) + \delta \geq H_n(\theta(X))$ holds for any $\delta > 0$, we have $H_n(X) \geq H_n(\theta(X))$. Also we have

$$\varphi_1(\mathfrak{U}_1, X) = \lim_{n \rightarrow \infty} \frac{1}{n} H_n(X) \geq \lim_{n \rightarrow \infty} \frac{1}{n} H_n(\theta(X)) = \varphi_2(\mathfrak{U}_2, \theta(X))$$

and also

$$\varphi_1(\mathfrak{U}_1) = \sup\{\varphi_1(\mathfrak{U}_1, X), X\} \geq \varphi_2(\mathfrak{U}_2, \theta(X)).$$

Suppose B is any partition of u_2 . Then $X = \theta^{-1}(B)$ is a partition of u_1 and $\theta(X) = B$. Therefore $\varphi_1(\mathfrak{U}_1) \geq \varphi_2(\mathfrak{U}_2, \theta(X)) = \varphi_2(\mathfrak{U}_2, B)$

for any B , hence $\varphi_1(\mathfrak{U}_1) \geq \varphi_2(\mathfrak{U}_2)$.

□



3. CONCLUSIONS

The study of the concept of entropy is very important in contemporary sciences. Entropy has been applied in information theory, physics, computer sciences, statistics, chemistry, biology, sociology, general systems theory and many other fields. In the paper, by using the concept of logical entropy of a partition in a product Łukasiewicz near semiring, we introduced the notion of logical entropy of a product Łukasiewicz near semiring dynamical systems. Also, we have defined a general type of entropy of a product MV-algebra dynamical system. Specifically, we introduced the notion of logical entropy of a product Łukasiewicz semiring and prove that entropy measure is invariant under isomorphism of product Łukasiewicz semiring dynamical systems.

REFERENCES

- [1] M. E. Anderson and T. H. Feil, *Lattice-Ordered Groups: An Introduction*, D. Reidel; U.S.A. and Canada: Kluwer Academic Publishers, Dordrecht, Holland, 1988.
- [2] L. P. Belluce, A. Di Nola, and A.R. Ferraioli, *MV-semirings and their sheaf representations*, *Order*, 30 (2013), 165–179.
- [3] S. Bonzio, I. Chajda, and A. Ledda, *Representing quantum structures as near semirings*, *Logic Journal of the IGPL*, 24 (2016), 719–742.
- [4] S. Bonzio, I. Chajda, and A. Ledda, *Representing quantum structures as near semi rings*, *Logic Journal of the IGPL*, 4 (2016), 719–742.
- [5] I. Chajda, *Algebraic theory of tolerance relations*, University Palack, 1991.
- [6] I. Chajda and H. Langer, *Commutative basic algebras and coupled near semirings*, *Soft Computing*, 19 (2015), 1129–1134.
- [7] M. Dellnitz and O. Junge, *Set oriented numerical methods for dynamical systems*, *Handbook of dynamical systems*, 2 (2002), 221–264.
- [8] G. Hummer and I. G. Kevrekidis, *molecular dynamics of a peptide fragment: Free energy, kinetics, and long-time dynamics computations*, *The Journal of chemical physics*, 118(23) (2003), 10762–10773.
- [9] P. Holmes, J. L. Lumley, G. Berkooz, and C. W. Wand Rowley, *Turbulence, coherent structures, dynamical systems and symmetry*, Cambridge University Press, 2012.
- [10] I. G. Kevrekidis, L. D. Schmidt, and R. Aris, *On the dynamics of periodically forced chemical reactors*, *Chemical engineering communications*, 30(6) (1984), 323–330.
- [11] A. N. Kolmogorov, *New metric invariant of transitive dynamical systems and automorphisms of Lebesgue spaces*, *Dokl. Russ. Acad. Sci.*, 119 (1958), 861–864.
- [12] W. M. Liu, H. W. Hethcote, and S. A. Levin, *Dynamical behavior of epidemiological models with nonlinear incidence rates*, *Journal of mathematical biology*, 25(4) (1987), 359–380.
- [13] I. Mezic, *Analysis of fluid flows via spectral properties of the Koopman operator*, *Annual Review of Fluid Mechanics*, 45 (2013), 357–378.
- [14] A. Molkhasi, *On strongly algebraically closed lattices*, *J Sib. Fed. Univ. Math. Phys.*, 9(2) (2016), 202–208.
- [15] A. Molkhasi, *On some strongly algebraically closed semirings*, *Journal of Intelligent and Fuzzy Systems*, 36(6) (2019), 6393–6400.
- [16] A. Molkhasi and K. P. Shum, *Strongly algebraically closed orthomodular near semirings*, *Rendiconti del Circolo Matematico di Palermo Series 2*, 69(3) (2020), 803–812.
- [17] D. Markechovi, A. Ebrahimzadeh, and Z. Eslami Giski, *Logical entropy of dynamical systems*, *Adv. Differ. Equ.*, 2018(70) 2018.
- [18] D. Markechovi and B. Riecan, *Logical entropy of dynamical systems in product MV-algebras and general scheme*, *Advances in Continuous and Discrete Models*, 9 (2019).
- [19] D. Mundici, *Interpretation of AFC*-algebras in Łukasiewicz sentential calculus*, *J. Funct. Anal.*, 56 (1986), 889–894.
- [20] H. Poincare, *Sur le probleme des trois corps et les equations de la dynamique*, *Acta Mathematica*, 13 (1890), A3–A270.



- [21] J. Petrovicovi, *On the entropy of partitions in product MV-algebras*, *Soft Comput.*, 4 (2000), 41–44.
- [22] J. Petrovicovi, *On the entropy of dynamical systems in product MV-algebras*, *Fuzzy Sets Syst.*, 121 (2001), 347–351.
- [23] B. Riecan and D. Markechovi, *φ -Entropy of IF-partitions*, *Notes IFS*, 23 (2017), 9–15.
- [24] S. H. Strogatz, *Nonlinear dynamics and chaos: with applications to physics, biology, chemistry, and engineering*, CRC press, 2018.
- [25] C.F. Shannon and W. Weaver, *Mathematical theory of communication*, *University of Illinois Press*, Urbana, 1949.
- [26] Y.G. Sinai, *On the notion of entropy of a dynamical system*, *Dokl. Russ. Acad. Sci.*, 124 (1959), 768–771.
- [27] J. Walleczek, *Self-organized biological dynamics and nonlinear control: toward understanding complexity, chaos and emergent function in living systems*, Cambridge University, 2006.

Uncorrected Proof

