



Application of new Kudryashov method to Sawada-Kotera and Kaup-Kupershmidt equations

Zeynep Aydn and Filiz Tacan

Faculty of Science, Department of Mathematics and Computer, Eskisehir Osmangazi University, Eskisehir, Turkey.

Abstract

In this article, with the help of the new Kudryashov method, we examine general solutions to the (2+1)-dimensional Sawada-Kotera equation (SKE) and Kaup-Kupershmidt (KK) equation. Using Maple, a symbolic computing application, it was shown that all obtained solutions are given by hyperbolic, exponential and logarithmic function solutions which obtained solutions are useful for fluid dynamics, optics and so on. Finally, we have presented some graphs for general solutions of these equations with special parameter values. The reliability and scope of programming provide eclectic applicability to high-dimensional nonlinear evolution equations for the development of this method. The results found gave us important information regarding the applicability of the new Kudryashov method.

Keywords. Exact solutions, Travelling wave solutions, Nonlinear partial differential equations.

2010 Mathematics Subject Classification. 65L05, 34K06, 34K28.

1. INTRODUCTION

Partial differential equations are important component of applied mathematics and have been known by scientists for a long time. They have a significant role in many fields including elasticity theory, fluid dynamics, plasma physics and engineering [22]. The development of computer programs for engineers and applied scientists approach to problems in PDEs has had an a signifiical impact. Pachage programs such as Maple, Mathematica, Matlab have started to be used for large calculations in the solutions of problems. In this paper, we used the symbolic computer program Maple to determine the exact solutions of partial differential equations.

Different approaches to solving nonlinear partial differential equations have been developed in recent years by scientits. Such as (G'/G) -expansion method [28], symmetry method [2], tanh-coth method [23, 30], Hirota's bilinear method [12], sine-cosine function method [32], the homogenous balance method [7, 29], simplest equation method [5, 34], Backlund transformation method [24], exp-function method [10], variational iteration method [11], sine-Gordon method [33], first integral method [8], P^6 - model expansion method [1], Kudryashov method [35], unified Ricatti equation expansion method [36], trial function method [3], sine-Gordon expansion approach [6], and so on.

Kudryashov method is one of the methods to solve the nonlinear PDEs. Kudryashov method was proposed by N. A. Kudryashov in 2011 [16–18]. Many researchers developed new methods using Kudryashov method [25, 37]. Like as modified Kudryashov method [14, 15, 27], improved Kudryashov method, modified improved Kudryashov method, generalized Kudryashov method, modified genereralized Kudryashov method, improved generalized Kudryashov method, modified improved generalized Kudryashov method, extended Kudryashov method and the new Kudryashov method [19] that we use in this article.

In this paper we will explain the new Kudryashov method proposed by N.A. Kudryashov. Then we will apply this method to the (2+1)-dimensional Sawada Kotera equation and the (1+1)-dimensional Kaup-Kuperchmidt equation. The new Kudryashov method is introduced in section 2. The exact solutions to the (2+1)-dimensional Sawada Kotera

Received: 09 January 2024 ; Accepted: 15 June 2024.

*Corresponding author. Email: zeynepaydinn10@gmail.com .

equation and the (1+1)-dimensional Kaup-Kupershmidt equation are the found and illustrated in section 3. Also, we present some visual representation of the solutions. The final section provided the conclusion.

2. THE NEW KUDRYASHOV METHOD

In this section, we describe the main steps of the new Kudryashov method. Assume the next nonlinear partial differential equation (PDE):

$$\Omega(u, u_t, u_x, u_{tt}, u_{xy}, u_{xx}, \dots) = 0, \tag{2.1}$$

where Ω simply represents a polynomial.

Step 1 By the travelling wave transformation $\vartheta = x + y - \sigma t$ and with $u(x, y, t) = U(\vartheta)$, Equation (2.1) can be reduced to an ordinary differential equation (ODE)

$$G(U, U_{\vartheta}, U_{\vartheta\vartheta}, U_{\vartheta\vartheta\vartheta}, \dots) = 0. \tag{2.2}$$

Step 2 Consider the Equation (2.2) has a solution as:

$$U(\vartheta) = \sum_{i=0}^N c_i Q^i(\vartheta), \tag{2.3}$$

where c_i ($i = 0, 1, \dots, N$) are the coefficients of $Q^i(\vartheta)$ with $c_N \neq 0$ and

$$Q(\vartheta) = \frac{1}{(aA^{(\Theta\vartheta)} + bA^{(-\Theta\vartheta)}),} \tag{2.4}$$

is the solution of the differential Equation (2.5)

$$\left(Q'(\vartheta)\right)^2 = (\Theta(\ln A)Q(\vartheta))^2(1 - 4abQ^2(\vartheta)), \tag{2.5}$$

where a, b, Θ and A are constants and non-zero arbitrary real parameters to be determined later, with $A > 0$ and $A \neq 0$. Also the positive integer N can be determined by using homogenous balance between the highest order derivative term with the highest order nonlinear term appearing in ODE (2.2).

Step 3 After substituting solution (2.3) into Equation (2.2), the left-hand side of Equation (2.2) can be converted into a polynomial in powers of $Q(\vartheta)$. Collecting the terms that include the same power of $Q(\vartheta)$ and equating each coefficient equal to zero, one obtains an algebraic equation system for c_0, c_1, c_2, A, a, b and Θ .

Step 4 We solve the algebraic equations in the Step 3 with the help of the Maple. Substituting the obtained values of c_0, c_1, c_2, A, a, b and Θ into solution (2.3) by considering Equation (2.4), the solutions of the NLPE in Equation (2.1) can be obtained.

3. APPLICATION OF THE NEW KUDRYASHOV METHOD

3.1. Solutions of the (2+1)-dimensional Sawada-Kotera Equation. One of the important model in mathematical physics is the classical Sawada-Kotera equation (SKE), which was put forth by Sawada and Kotera firstly [32]. Application for this equation include nonlinear optics, nonlinear acoustic waves in an inharmonic lattice and quantum mechanics. It also emerged to be used to describe how long waves move in shallow water [33, 34]. Numerous varieties of approximate and exact solutions are obtained, such as lump solutions, periodic solitary wave solutions, N soliton solutions, and two soliton solutions. Thus, investigating novel interaction solutions among soliton molecules, breather molecules and soliton-breather molecules of it also of great significance.

The following form of the (2+1)-dimensional Sawada-Kotera Equation was proposed by Ma et al. [21].

$$u_t + u_{xxy} - 3uu_y - 3u_xv + u_{5x} - 15(uu_{xx} - u^3)_x = 0, \text{ with } v_x = u_y, \tag{3.1}$$

where $u = u(x, y, t)$ and $v = v(x, y, t)$ are the unknow function with a real value that results from the real independent variables x, y and t . Also x, y are dimensionless spatial variables and t is temporal variable.

Now, we apply the new Kudryashov method to find the exact-solution for the Equations (3.1).



Using the travelling wave transformation

$$u(x, y, t) = U(\vartheta), v(x, y, t) = V(\vartheta), \vartheta = x + y - \sigma t. \quad (3.2)$$

where σ is represents the wave velocity. The Equation (3.1) is carried to NODE

$$-\sigma U_{\vartheta} + U_{\vartheta\vartheta\vartheta} - 3UU_{\vartheta} - 3U_{\vartheta}V + U_{\vartheta\vartheta\vartheta\vartheta} - 15(UU_{\vartheta\vartheta} - U^3)_{\vartheta} = 0. \quad (3.3)$$

$$U_{\vartheta} = V_{\vartheta}. \quad (3.4)$$

By applying integrate on Equation (3.4) with respect to ϑ and then taking the integral constant as zero, we get

$$U = V. \quad (3.5)$$

Substituting Equation (3.5) into Equation (3.3), after integrating with respect to ϑ ,

$$-\sigma U + U_{\vartheta\vartheta} - 3U^2 + U_{\vartheta\vartheta\vartheta} - 15(UU_{\vartheta\vartheta} - U^3) + B = 0. \quad (3.6)$$

where B is constant of integration.

By balancing the highest order derivative term $U_{\vartheta\vartheta\vartheta}$ and the highest order nonlinear term U^3 , we get;

$$N + 4 = 3N, \quad (3.7)$$

$$N = 2.$$

So from solution (2.3), we can write

$$U(\vartheta) = c_0 + c_1 Q(\vartheta) + c_2 Q^2(\vartheta), \quad (3.8)$$

where c_0, c_1 and c_2 are constants to be determined later.

Substituting solution (3.8) into the Equation (3.6) yields a polinomial in $Q(\vartheta)$. A system of algebraic equations is obtained by setting each coefficient of the equations to zero.

$$Q^6 : 1920 \ln(A)^4 a^2 b^2 c_2 \Theta^4 + 360 \ln(A)^2 abc_2^2 \Theta^2 + 15c_2^3,$$

$$Q^5 : 384 \ln(A)^4 a^2 b^2 c_1 \Theta^4 + 480 \ln(A)^2 abc_1 c_2 \Theta^2 + 45c_1 c_2^2,$$

$$Q^4 : -480 \ln(A)^4 abc_2 \Theta^4 + 360 \ln(A)^2 abc_0 c_2 \Theta^2 + 120 \ln(A)^2 abc_1^2 \Theta^2 - 24 \ln(A)^2 abc_2 \Theta^2 - 60 \ln(A)^2 c_2^2 \Theta^2 + 45c_0 c_2^2 + 45c_1^2 c_2 - 3c_2^2,$$

$$Q^3 : -80 \ln(A)^4 abc_1 \Theta^4 + 120 \ln(A)^2 abc_0 c_1 \Theta^2 - 8 \ln(A)^2 abc_1 \Theta^2 - 75 \ln(A)^2 c_1 c_2 \Theta^2 + 90c_0 c_1 c_2 + 15c_1^3 - 6c_1 c_2,$$

$$Q^2 : 16 \ln(A)^4 \Theta^4 c_2 - 60 \ln(A)^2 \Theta^2 c_0 c_2 - 15 \ln(A)^2 \Theta^2 c_1^2 + 4 \ln(A)^2 \Theta^2 c_2 + 45c_0^2 c_2 + 45c_1^2 c_0 - \sigma c_2 - 6c_0 c_2 - 3c_1^2,$$

$$Q^1 : \ln(A)^4 \Theta^4 c_1 - 15 \ln(A)^2 \Theta^2 c_0 c_1 + \ln(A)^2 \Theta^2 c_1 + 45c_0^2 c_1 - \sigma c_1 - 6c_0 c_1,$$

$$Q^0 : 15c_0^3 - \sigma c_0 - 3c_0^2 + B.$$

To solve the above system of algebraic equations we use Maple Software. Therefore we get following 2 solution sets;

Set 1

$$\begin{aligned} B &= 16 \ln(A)^4 \Theta^4 c_0 - 60 \ln(A)^2 \Theta^2 c_0^2 + 4 \ln(A)^2 \Theta^2 c_0 + 30c_0^3 - 3c_0^2, \\ \sigma &= 16 \ln(A)^4 \Theta^4 - 60 \ln(A)^2 \Theta^2 c_0 + 4 \ln(A)^2 \Theta^2 + 45c_0^2 - 6c_0, \\ c_0 &= c_0, c_1 = 0, c_2 = -8 \ln(A)^2 \Theta^2 ab. \end{aligned} \quad (3.9)$$

Substituting Equations (3.9) into (3.8) with using (2.4), we obtain the exact solution of the (2+1)-dimensional SKE as follows:

$$U_1 = V_1 = c_0 - \frac{8 \ln(A)^2 \Theta^2 ab}{(aA^{(\Theta\vartheta)} + bA^{(-\Theta\vartheta)})^2}, \quad (3.10)$$

where

$$\vartheta = - \left(16 \ln(A)^4 \Theta^4 - 60 \ln(A)^2 \Theta^2 c_0 + 4 \ln(A)^2 \Theta^2 + 45c_0^2 - 6c_0 \right) t + x + y. \quad (3.11)$$



Set 2

$$B = \frac{-(128 \ln(A)^6 \Theta^6)}{9} + \frac{(16 \ln(A)^4 \Theta^4)}{15} - \frac{1}{225}, \sigma = 16 \ln(A)^4 \Theta^4 - \frac{1}{5},$$

$$c_0 = \frac{(4 \ln(A)^2 \Theta^2)}{3} + \frac{1}{5}, c_1 = 0, c_2 = -16 \ln(A)^2 \Theta^2 ab.$$
(3.12)

Substituting Equations (3.12) into (3.8) with using ((2.4), we obtain exact solution of the (2+1)-dimensional SKE as follows:

$$U_2 = V_2 = \frac{(4 \ln(A)^2 \Theta^2)}{3} + \frac{1}{15} - \frac{16 \ln(A)^2 \Theta^2 ab}{(aA^{(\Theta\vartheta)} + bA^{(-\Theta\vartheta)})^2},$$
(3.13)

where

$$\vartheta = -\left(16 \ln(A)^4 \Theta^4 - \frac{1}{5}\right)t + x + y.$$
(3.14)

In particular, if we have $a = b$ in solution U_2 , then Equation (3.3) has the following bright-soliton solution

$$u_{2.1}(x, y, t) = \frac{1}{15} + \frac{4 \ln(A)^2 \Theta^2 \left(1 - 3 \operatorname{sech}^2 \left(\Theta \left(80t \ln(A)^4 \Theta^4 - t - 5x - 5y\right) \frac{\ln(A)}{5}\right)\right)}{3}.$$
(3.15)

On the other hand if we take $a = -b$ in in solution U_2 , then Equation (3.3) has the following singular-soliton solution

$$u_{2.2}(x, y, t) = \frac{4 \ln(A)^2 \Theta^2}{3} + 4 \ln(A)^2 \Theta^2 \operatorname{csch}^2 \left(\Theta \left(80t \ln(A)^4 \Theta^4 - t - 5x - 5y\right) \frac{\ln(A)}{5}\right) + \frac{1}{15}.$$
(3.16)

Figure 1 and Figure 2 as given below, eight graphs in total are drawn with the help of the Maple for the exact solutions $U_1 = V_1$ (3.10) and $U_2 = V_2$ (3.13). For the graphs of these exact solutions are drawn for the Sawada-Kotera equation, the parameters and the range of x and t values are specially chosen.

3.2. Solutions of the (1+1)-dimensional Kaup-Kupershmidt (KK) equation. The standard fifth-order KdV equation (fKdV) of the form [31],

$$u_t + \alpha u^2 u_x + \beta u_x u_{xx} + \gamma u u_{3x} + u_{5x} = 0.$$
(3.17)

where α, β and γ are arbitrary nonzero and real parameters and $u = u(x, t)$ is a differentiable function. With taking different values of the parameters α, β and γ , the Equation (3.17) change the characteristics of the fKdV equation.

Kaup-Kupershmidt equation is characterized by

$$\beta = \frac{5}{2}\gamma, \alpha = \frac{1}{5}\gamma^2.$$

For $\gamma = 10$, then $\beta = 25$ and $\alpha = 20$, therefore Equation (3.17) reduces to standard KK equation:

$$u_t + 20u^2 u_x + 25u_x u_{xx} + 10u u_{3x} + u_{5x} = 0.$$
(3.18)

where $u = u(x, t)$ is the unknow function with a real value that results from the real independent variables x and y .

The Kaup-Kupershmidt equation was proposed by Kaup in 1980 [13, 20].

To find the exact-solution for the Kaup-Kupershmidt Equation (3.18), we apply the new Kudryashov method.

Consider the following travelling wave transformation

$$u(x, t) = U(\vartheta), \vartheta = x - \sigma t.$$
(3.19)

where σ is represents the wave velocity.

The Equation (3.18) is carried to NODE

$$-\sigma U_\vartheta + 20U^2 U_\vartheta + 25U_\vartheta U_{\vartheta\vartheta} + 10U_\vartheta U_{\vartheta\vartheta\vartheta} + U_{5\vartheta} = 0,$$
(3.20)






FIGURE 1. New exact solution $U_1 = V_1$ (3.6) of the (2+1)-dimensional SKE (3.10), for $A = 1.5, \Theta = 1.5, a = 3, b = 2, c_0 = 0.5, y = 1$, when in the range of $t \in [0, 10], x \in [-10, 10]$.

integrating with respect to ϑ

$$-\sigma U + \frac{20}{3}U^3 + \frac{15}{2}(U_{\vartheta})^2 + 10UU_{\vartheta\vartheta} + U_{\vartheta\vartheta\vartheta} + B = 0, \tag{3.21}$$

where B is constant of integration. By balancing the highest order derivative term $U_{\vartheta\vartheta\vartheta}$ and the highest order nonlinear term U^3 , we get,

$$\begin{aligned} N + 4 &= 3N, \\ N &= 2. \end{aligned} \tag{3.22}$$

So from (2.3), we can write

$$U(\vartheta) = c_0 + c_1Q(\vartheta) + c_2Q^2(\vartheta), \tag{3.23}$$

where c_0, c_1 and c_2 are constants to be determined later.

Substituting (3.23) into the Equation (3.21) yields a polinomial in $Q(\vartheta)$. A system of algebraic equations is obtained by setting each coefficient of the equations to zero.

$$\begin{aligned} Q^6 : & 1920 \ln(A)^4 \Theta^4 a^2 b^2 c_2 - 360 \Theta^2 \ln(A)^2 abc_2^2 + \frac{(20c_2^3)}{3}, \\ Q^5 : & 384 \ln(A)^4 \Theta^4 a^2 b^2 c_1 - 440 \Theta^2 \ln(A)^2 abc_1 c_2 + 20c_1 c_2^2, \\ Q^4 : & -480 \ln(A)^4 \Theta^4 abc_2 - 240 \Theta^2 \ln(A)^2 abc_0 c_2 - 110 \Theta^2 \ln(A)^2 abc_1^2 + 70 \Theta^2 \ln(A)^2 c_2^2 + 20c_0 c_2^2 + 20c_1^2 c_2, \end{aligned}$$






FIGURE 2. New exact solution $U_2 = V_2(3.13)$ of the (2+1)-dimensional SKE (3.6), for $A = 1.5, \Theta = 2, a = 5, b = 2, c_0 = 0.5, y = 2$ when in the range of $t \in [0, 10], x \in [-10, 10]$.

$$\begin{aligned}
 Q^3 &: 80\Theta^2 \ln(A)^2 c_1 c_2 - 80 \ln(A)^4 \Theta^4 abc_1 + 40c_0 c_1 c_2 + \frac{20c_1^3}{3} - 80\Theta^2 \ln(A)^2 abc_0 c_1, \\
 Q^2 &: 40\Theta^2 \ln(A)^2 c_0 c_2 + 20c_0 c_1^2 - \sigma c_2 + 20c_0^2 c_2 + 16 \ln(A)^4 \Theta^4 c_2 + \frac{35\Theta^2 \ln(A)^2 c_1^2}{2}, \\
 Q^1 &: \ln(A)^4 \Theta^4 c_1 + 10\Theta^2 \ln(A)^2 c_0 c_1 + 20c_0^2 c_1 - \sigma c_1, \\
 Q^0 &: B - \sigma c_0 + \frac{20c_0^3}{3}.
 \end{aligned}$$

To solve the above system of algebraic equations, we use Maple Software and we get the following two solution sets; **Set 1**

$$B = \frac{\Theta^6 \ln(A)^6}{3}, \quad \sigma = \ln(A)^4 \Theta^4, \quad c_0 = \frac{-\Theta^2 \ln(A)^2}{2}, \quad c_1 = 0, \quad c_2 = 6\Theta^2 ab \ln(A)^2. \tag{3.24}$$

Substituting Equations (3.24) into (3.23) using (2.4), we obtain the exact solution of the KK as follows:

$$U_3 = \frac{-\Theta^2 \ln(A)^2}{2} + \frac{6 \ln(A)^2 \Theta^2 ab}{(aA^{(\Theta\vartheta)} + bA^{(-\Theta\vartheta)})^2}, \tag{3.25}$$

where

$$\vartheta = -\ln(A)^4 \Theta^4 t + x. \tag{3.26}$$






FIGURE 3. New exact solution $U_3(3.25)$ of the KK Equation (3.21), for $A = 2.5, \Theta = 1.5, a = 1, b = 2$ when in the range of $t \in [0, 10], x \in [-10, 10]$.

In particular, if we have $a = b$ in solution U_2 , then Equation (3.3) has the following bright-soliton solution

$$u_{1.1}(x, t) = \Theta^2 \ln(A)^2 \frac{3 \operatorname{sech}^2 \left(\Theta^2 \left(\ln(A)^4 \Theta^8 t - x \right) \ln(A) \right) - 1}{2}. \tag{3.27}$$

On the other hand if we take $a = -b$ in in solution U_2 , then Equation (3.3) has the following singular-soliton solution

$$u_{1.2}(x, t) = -\Theta^2 \ln(A)^2 \frac{1 + 3 \operatorname{csch}^2 \left(\Theta \left(\ln(A)^4 \Theta^4 t - x \right) \ln(A) \right)}{2}. \tag{3.28}$$

Set 2

$$B = \frac{-832\Theta^6 \ln(A)^6}{3}, \sigma = 176 \ln(A)^4 \Theta^4, c_0 = -4\Theta^2 \ln(A)^2, c_1 = 0, c_2 = 48\Theta^2 ab \ln(A)^2. \tag{3.29}$$

Substituting Equations (3.29) into (3.23) with using (2.4), we obtain the exact solution of the KK as follows:

$$U_4 = -4\Theta^2 \ln(A)^2 + \frac{48 \ln(A)^2 \Theta^2 ab}{(aA^{(\Theta\vartheta)} + bA^{(-\Theta\vartheta)})^2}, \tag{3.30}$$

where

$$\vartheta = -176 \ln(A)^4 \Theta^4 t + x. \tag{3.31}$$

The graphs are drawn with help of the Maple for the U_3 (3.25) and U_4 (3.30) by choosing the parameters and x, t values in Figures 3 and 4.






FIGURE 4. New exact solution $U_4(3.30)$ of the KK Equation (3.21), for $A = 3, \Theta = 0.2, a = 2, b = 5$ when in the range of $t \in [0, 10], x \in [-10, 10]$.

4. CONCLUSION

In this study, we discussed (2+1)-dimensional Sawada Kotera equation and the (1+1)-dimensional Kaup-Kupershmidt equation, both of which have several uses in mathematics and physics. We utilized the new Kudryashov method after reducing the PDE to ODE using traveling wave transformation. We observed different bright soliton solutions for the (2+1)-dimensional Sawada Kotera equation and the (1+1)-dimensional Kaup-Kupershmidt equation using the $Q(\vartheta)$ function, which is in exponential form and is the solution of the auxiliary equation. Various three and two dimensional plots, polar coordinate plots and contour plots, and dynamical characteristics of these waves are shown well using Maple. The newly found hyperbolic function solutions may be useful for understanding physical phenomena especially in long-wave propagation, dynamics of shallow water wave, and plasma fluid. Additionally, it is clear that the approach used in this article is a powerful and practical mathematical tool that may be used to provide exact solutions to various other different NLPDEs in the future.



REFERENCES

- [1] A. H. Bhrawy, et al., *Dispersive optical solitons with Schrödinger-Hirota equation.*, Journal of Nonlinear Optical Physics & Materials, *23*(01) (2014), 1450014.
- [2] G. W. Bluman and S. Kumei, *Symmetries and Differential Equations*, Springer-Verlag, New York, 1989.
- [3] Y. Chen, et al., *Dark wave, rogue wave and perturbation solutions of Ivancevic option pricing model.*, Nonlinear Dynamics, *105* (2021), 2539–2548.
- [4] K. Debin, H. Rezazadeh, N. Ullah, J. Vahidi, U. T. Kalim, and L. Akinyemi, *New soliton wave solutions of a (2+1)-dimensional Sawada-Kotera equation*, Journal of Ocean Engineering and Science, *8* (2023), 527–532.
- [5] M. Eslami, M. Mirzazadeh, and A. Biswas, *Soliton solutions of the resonant nonlinear Schrödinger equation in optical fibers with time dependent coefficients by simplest equation approach*, J. Mod. Opt., *60*(19) (2013), 16271636.
- [6] M. R. A. Fahim, et al., *Wave profile analysis of a couple of (3+ 1)-dimensional nonlinear evolution equations by sine-Gordon expansion approach.* Journal of Ocean Engineering and Science, *7*(3) (2022), 272279.
- [7] E. Fan and H. Zhang, *A note on the homogeneous balance method*, Phys. Lett., A *246* (1998), 403406.
- [8] Z. S. Feng, *The First integral method to study the Burgers-KdV equation*, J. Phys. A: Math. Gen., *35* (2002), 343349.
- [9] A. K. Gupta and S. S. Ray., *Numerical treatment for the solution of fractional fifth-order Sawada-Kotera equation using second kind Chebyshev wavelet method* Applied Mathematical Modelling, *39* (2015), 51215130.
- [10] J. H. He and X. H. Wu, *Exp-function method for nonlinear wave equations*, Chaos, Solitons & Fractals, *30*(3) (2006), 700708.
- [11] J. H. He, *Approximate solution of nonlinear differential equations with convolution product nonlinearities*, Computer Methods in Applied Mechanics and Engineering, *167*(1-2) (1998), 6973.
- [12] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge University Press, Cambridge, 2004.
- [13] D. Kaup, *On the inverse scattering problem for the cubic eigenvalue problems of the class $\psi_3x + 6Q\psi_x + 6R\psi = \lambda\psi$* , Stud. Appl. Math., *62* (1980), 189216.
- [14] A. Khaled, *The Modified Kudryashov Method for Solving Some Seventh Order Nonlinear PDEs*, Mathematical Physics, *11* (2015), 308319.
- [15] A. Kilicman and R. Silambarasan, *Modified Kudryashov method to solve generalized Kuramoto-Sivashinsky equation*, Symmetry *10*, *10* (2018), 527.
- [16] N. A. Kudryashov, *One method for finding exact solutions of nonlinear differential equations*, Commun. Nonlinear Sci. Numer. Simul. Commun. Nonlinear Sci. Number, *17* (2011).
- [17] N. A. Kudryashov, *Method of the logistic function for finding analytical solutions of nonlinear differential equations*, Model. Anal. Inf. Syst., *22* (2015), 2337.
- [18] N. A. Kudryashov, *Logistic function as solution of many nonlinear differential equations*, Appl. Math. Model., *39*(18) (2015), 57335742.
- [19] N. A. Kudryashov, *Method for finding highly dispersive optical solitons of nonlinear differential equations*, Optik, *206* (2020), 63550.
- [20] B. A. Kupershmidt, *A super KdV equation: an integrable system*, Phys. Lett. A, *102* (1984), 213215.
- [21] W. X. Ma, R. Zhou, and L. Gao, *Exact one-periodic and two-periodic wave solutions to Hirota bilinear equations in (2+1) dimensions.*, Modern Physics Letters A, *24*(21) (2009), 16771688.
- [22] C. L. Mader, *Numerical Modeling of Water Waves*, Numerical Modeling of Water Waves, Jun. 2004.
- [23] W. Malfliet and W. Hereman, *The tanh method. I: exact solutions of nonlinear evolution and wave equations*, Phys. Scripta, *54* (1996), 563568.
- [24] R. M. Miura, *Backlund Transformation*, Springer-Verlag, New York, 1973.
- [25] M. Ozisik, A. Secer, M. Bayram, and H. Aydin, *An encyclopedia of Kudryashovs integrability approaches applicable to optoelectronic devices*, Optik, *265* (2022), 169499.
- [26] K. Sawada and T. Kotera, *A Method Finding N-Soliton Solutions of the Kdv and KdV-Like Equations*, Prog. Theor. Phys., *51* (1974), 135567.



- [27] H. Srivastava, D. Baleanu, M. T. M. Jos, H. Rezazadeh, S. Arshed, and G. Hatira, *Traveling wave solutions to nonlinear directional couplers by modified Kudryashov method*, Phys. Scr., 95 (2020), 075217.
- [28] M. Wang, X. Li, and J. Zhang, *The (G/G) -expansion method and travelling wave solutions of nonlinear evolution equations in mathematical physics*, Phys. Lett. A., 372 (2008), 417423.
- [29] M. L. Wang, *Solitary wave solutions for variant Boussinesq equations*, Phys. Lett. A, 199 (1995), 169172.
- [30] A. M. Wazwaz, *The tanhcoth method for solitons and kink solutions for nonlinear parabolic equations*, Appl. Math. Comput. 188 (2007), 14671475.
- [31] A. M. Wazwaz, *Abundant solitons solutions for several forms of the fifth-order KdV equation by using the tanh method*, Appl. Math. Comput., 182 (2006), 283300.
- [32] C. Yan, *A simple transformation for nonlinear waves*, Phys. Lett. A., 224 (1996), 7784.
- [33] C. Yan, *A simple transformation for nonlinear waves*, Physics Letters A, 224(1-2) (1996), 7784.
- [34] A. Yildirim, A. S. Paghaleh, M. Mirzazadeh, and H. Moosaei, *A. Biswas, New exact traveling wave solutions for DS-I and DS- II equations*, Nonlinear Anal.: Modell. Control. 17(3) (2012), 369378.
- [35] E. M. E. Zayed , R. M. A. Shohib, and M. E. M. Alngar, *Cubic-quartic optical solitons in Bragg gratings fibers for NLSE having parabolic non-local law nonlinearity using two integration schemes*, Optical and Quantum Electronics, 53(8) (2021), 452.
- [36] E. M. E. Zayed, M. E. M. Alngar, and R. M. A. Shohib, *Cubic-quartic embedded solitons with χ (2) and χ (3) nonlinear susceptibilities having multiplicative white noise via It calculus*, Chaos, Solitons & Fractals, 168 (2023), 113186.
- [37] E. M. E. Zayed, et al. *Solitons in magneto-optics waveguides for the nonlinear BiswasMilovic equation with Kudryashovs law of refractive index using the unified auxiliary equation method*. Optik, 235 (2021), 166602.

