



Hyers-Ulam and exponential stabilities of autonomous and non-autonomous difference equations

Gul Rahmat¹, Afaq Ahmad¹, Muhammad Sarwar^{2,3}, Kamaleldin Abodayeh³, Cemil Tunç^{4,*}

¹Department of Mathematics, Islamia College Peshawar, Pakistan.

²Department of Mathematics, University of Malakand, Dir Lower, Pakistan.

³Department of Mathematics, Department of Mathematics, Prince Sultan University, Saudi Arabia.

⁴Department of Mathematics, Faculty of Sciences, Van Yuzuncu Yil University, Campus, 65080 Van, Turkey.

Abstract

In this manuscript, we studied the Hyers-Ulam and exponential stabilities of autonomous and non-autonomous difference equations of first and second order. At the end we provide some examples to support the results.

Keywords. Hyer-Ulam, Exponential Stability, Difference Equations.

2010 Mathematics Subject Classification. 35B35.

1. INTRODUCTION

Differential equations have been used to analyze a variety of biological systems and models for a very long time in mathematics. Differential equations have several uses in a variety of disciplines, including engineering, economics, statistics, and the natural sciences [1–4]. Even while differential equations are quite useful, some real-world issues call for discrete data sampling, which necessitates the usage of difference equations [5, 6]. Difference equations have become crucial tools in the study of biological systems as well as in the sciences, technology and mathematical physics. They offer a structure for precisely describing phenomena, such as wind flow dynamics and monographs, which helps in problem-solving in these fields.

Stability is one of the key qualitative characteristics of different systems that play a vital impact. System performance must be ensured by stability assessment. Due to its applicability in a variety of situations, Hyers-Ulam stability has attracted a lot of attention from researchers. The concept of Hyers-Ulam stability emerged almost eighty years ago in 1940 when Ulam [7] put a question before mathematicians in a conference about the existence of a group homomorphism which can be near to an approximate group homomorphism on groups. After a year in 1941, Hyers [8] addressed Ulam's puzzles by considering the group as a Banach space and provided solutions. Consequently, the stability was named Hyers-Ulam stability. Donald H. Hyers made significant contributions in this field by providing a partially positive response to Ulam's query in the case of additive mappings within Banach spaces. Since then, many works have been published regarding various extensions of Ulam's problem and Hyers' theorem. In 1978, Rassias [9] provided an overview of this stability and applied the concept to the Cauchy problem. Furthermore, Obloza [10] use this concept in the field of differential equations, while Jung [11] and Khan et al, [12] discussed the idea for difference equations. Z. Gao et al. [13] and S.O. Shah et al, [14] reported the same stability results on delayed first-order nonlinear dynamic system. Recent studies have focused on generalized Hyers-Ulam stability. A.R. Aruldass [15]. Additionally, investigations were made on the Hyers-Ulam stability of differential equations of order two utilizing the Mahgoub type. For other models and its solution we refer to [22] and [23]. Hyers-Ulam stability, is a mathematical concept related to functional equations. Hyers-Ulam stability is a mathematical concept that deals with the stability of functional equations. More precisely, it concerns the question of whether small perturbations of the inputs of a

Received: 23 December 2023 ; Accepted: 19 August 2024.

* Corresponding author. Email: cemtunc@yahoo.com .

functional equation lead to correspondingly small perturbations of the output. Additionally, Ulam type stabilities several different mathematical models have been invstigated [25–30] and some intereting result have been obtained in these sources.

In this paper, we describe the exponential stability and Hyers-Ulam stability of autonomous and non-autonomous difference systems of the type.

$$\begin{cases} y_{n+1} = (1+r)y_n, & n \geq 0, \\ y_0 = \theta, \end{cases} \quad (1.1)$$

$$\begin{cases} y_{n+2} = (1+r)y_n, & n \geq 0, \\ y_0 = a, \\ y_1 = b, \end{cases} \quad (1.2)$$

$$\begin{cases} y_{n+1} = r_n y_n, & n \geq 0, \\ y_0 = \theta, \end{cases} \quad (1.3)$$

and

$$\begin{cases} y_{n+2} = r_n y_n, & n \geq 0, \\ y_0 = \alpha, \\ y_1 = \beta, \end{cases} \quad (1.4)$$

where $r, r_n \in \mathbb{R}$, $y_n \in \mathbb{C}(\mathbb{Z}_+, \mathbf{X})$ where $\mathbb{C}(\mathbb{Z}_+, \mathbf{X})$ is the space of convergent sequences equipped with norm supremum, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ and $\mathbf{X} = \mathbb{R}^m$.

For other recent work related to the Hyers-Ulam stability of different type of difference equations we refer to [18], [19], [20], [21], and [24].

2. PRELIMINARIES

Throughout the paper we will use the following notations: the vector norm will be denoted by $\|\cdot\|$, the n -dimensional Euclidean space will be represented by the symbol $\mathbf{X} = \mathbb{R}^m$. The collection of real, integer, and non-negative integer numbers are represented by the symbols \mathbb{R}, \mathbb{Z} and \mathbb{Z}_+ , respectively.

Definition 2.1. We recall that a family $\mathcal{E} = \{E(n, k) : n \geq k \geq 0\}$ of bounded linear operators acting on a Banach space \mathbf{X} is called a discrete evolution family if it satisfy the following two conditions:

(i) $E(n, n) = I$ for all $n \geq 0$, (ii) $E(n, s)E(s, k) = E(n, k)$ for all $n \geq s \geq k \geq 0$.

In the solutions of non-autonomous systems we can take the evolution family as given below:

$$E(n, k) := \begin{cases} I, & \text{if all } n = k; \\ r_{n-1}r_{n-2}r_{n-3}\dots r_k, & \text{for all } n \geq k \geq 0. \end{cases}$$

Definition 2.2. The systems (1.1) to (1.4) are said to be uniformly exponentially stables if there are two positive real numbers ξ and μ , such that $\|y_n\| \leq \xi e^{-\mu n}$, $\forall n \geq 0$.

Definition 2.3. A sequence Ψ_n is referred to as an ϵ -approximate solution of the (1.1), (1.2), (1.3), and (1.4) if the following inequalities holds.

$$\|\Psi_{n+1} - (1+r)\Psi_n\| \leq \epsilon, \quad n \geq 0, \quad (2.1)$$

$$\|\Psi_{n+2} - (1+r)\Psi_n\| \leq \epsilon, \quad n \geq 0, \quad (2.2)$$



$$\|\Psi_{n+1} - r_n \Psi_n\| \leq \epsilon, \quad n \geq 0, \tag{2.3}$$

$$\|\Psi_{n+2} - r_n \Psi_n\| \leq \epsilon, \quad n \geq 0. \tag{2.4}$$

Remark 2.4. From Definition 2.3 we have that $y_n \in \mathbb{C}(\mathbb{Z}_+, \mathbf{X})$ will satisfies (2.1), (2.2), (2.3), and (2.4) if and only if there is a sequence $f \in \mathbb{C}(\mathbb{Z}_+, \mathbf{X})$ with $\|f_n\| \leq \epsilon$ such that

$$\begin{cases} y_{n+1} = (1+r)y_n + f_n, & n \geq 0, \\ y_0 = \theta, \end{cases}$$

$$\begin{cases} y_{n+2} = (1+r)y_n + f_n, & n \geq 0, \\ y_0 = a \\ y_1 = b, \end{cases}$$

$$\begin{cases} y_{n+1} = r_n y_n + f_n, & n \geq 0, \\ y_0 = \theta, \end{cases}$$

$$\begin{cases} y_{n+2} = r_n y_n + f_n, & n \geq 0, \\ y_0 = \alpha, \\ y_1 = \beta. \end{cases}$$

Lemma 2.5. The systems (1.1), (1.2), (1.3), and (1.4) have the solutions:

$$\begin{aligned} y_n &= (1+r)^n \theta, \\ y_n &= \begin{cases} (1+r)^{\frac{n}{2}} a, & \text{when } n \text{ is even,} \\ (1+r)^{\frac{n-1}{2}} b, & \text{when } n \text{ is odd,} \end{cases} \\ y_n &= E(n, 0)\theta, \end{aligned}$$

and

$$y_n = \begin{cases} \prod_{\lambda=1}^{\frac{n}{2}} r_{n-2\lambda} \alpha, & \text{when } n \text{ is even,} \\ \prod_{\lambda=1}^{\frac{n-1}{2}} r_{n-2\lambda} \beta, & \text{when } n \text{ is odd.} \end{cases}$$

respectively.

Lemma 2.6. The systems (1.1), (1.2), (1.3), and (1.4) have the approximate solutions:

$$\begin{aligned} \varphi_n &= (1+r)^n \gamma + \sum_{\lambda=0}^{n-1} (1+r)^{n-1-k} f_\lambda, \\ \psi_n &= \begin{cases} (1+r)^{\frac{n}{2}} a + \sum_{\lambda=0}^{\frac{n}{2}-1} (1+r)^{\frac{n}{2}-1-\lambda} f_{2\lambda}, & n = 2, 4, 6, 8, \dots \\ (1+r)^{\frac{n-1}{2}} b + \sum_{\lambda=0}^{\frac{n-1}{2}-1} (1+r)^{\frac{n-1}{2}-1-\lambda} f_{2\lambda+1}, & n = 1, 3, 5, 7, \dots \end{cases} \\ \phi_n &= E(n, 0)\theta + \sum_{\lambda=1}^n E(n, \lambda) f_{\lambda-1}, \end{aligned}$$

and

$$\Psi_n = \begin{cases} \prod_{\lambda=1}^{\frac{n}{2}} (r_{n-2\lambda}) \alpha + \sum_{j=1}^{\frac{n}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda}) f_{n-2\lambda-2} + f_{n-2}, & n = 2, 4, 6, 8, \dots \\ \prod_{\lambda=1}^{\frac{n-1}{2}} (r_{n-2\lambda}) \beta + \sum_{j=1}^{\frac{n-1}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda}) f_{n-2\lambda-2} + f_{n-2}, & n = 1, 3, 5, 7, \dots \end{cases}$$

respectively.

The solutions in Lemmas 2.5 and 2.6 can easily be obtained by putting the values of n .



Definition 2.7. The systems (1.1), (1.2), (1.3) and (1.4) are said to be Hyers-Ulam stable if there exists a positive real numbers \mathbf{L} such that for every approximate solutions ψ_n of the system (1.1), (1.2), (1.3) and (1.4) there are solutions y_n of (1.1), (1.2), (1.3) and (1.4) such that

$$\|y_n - \psi_n\| \leq \mathbf{L}\epsilon, \quad n \in I.$$

Remark 2.8. [17] Every semigroup is an evolution but the converse is not true in general.

Remark 2.9. [17] An evolution family is a semigroups if it is periodic of period one.

From Remarks 2.8 and 2.9 the relation between autonomous and non-autonomous is very clear that every autonomous may be consider as non-autonomus but for the reverse it is necessary that the time dependent matrix must be periodic with period 1.

3. HYERS-ULAM STABILITY OF AUTONOMOUS SYSTEMS

In this section we study the Hyers-Ulam stability of autonomous difference systems of first and second order. For both the results we need the following assumption:

$$\Lambda_1 : |1 + r| < 1.$$

Theorem 3.1. If Λ_1 holds then, the system (1.1) is Hyer-Ulam stable.

Proof. Since by Lemmas 2.5 and 2.6 the system (1.1) has the exact and approximate solutions:

$$y_n = (1 + r)^n \gamma$$

and

$$\varphi_n = (1 + r)^n \gamma + \sum_{\lambda=0}^{n-1} (1 + r)^{n-1-k} f_\lambda,$$

respectively.

Now consider

$$\begin{aligned} \|y_n - \varphi_n\| &= \|(1 + r)^n \gamma - (1 + r)^n \gamma - \sum_{\lambda=0}^{n-1} (1 + r)^{n-1-\lambda} f_\lambda\| \\ &= \left\| \sum_{\lambda=0}^{n-1} (1 + r)^{n-1-\lambda} f_\lambda \right\| \leq \epsilon \sum_{\lambda=0}^{n-1} (1 + r)^{n-1-\lambda} \\ &= \epsilon (a^{n-1} + a^{n-2} + \dots + a^0) = \epsilon (1 + a^1 + a^2 + \dots + a^{n-1}) \\ &\leq \epsilon (1 + a^1 + a^2 + \dots) = \frac{\epsilon}{1 - a} = \frac{\epsilon}{1 - (1 + r)} = \frac{\epsilon}{-r} = L\epsilon, \end{aligned}$$

where $L = \frac{1}{-r}$, hence the system is Hyers-Ulam stable. □

Theorem 3.2. If Λ_1 holds then, the system (1.2) is Hyer-Ulam stable.

Proof. Since by Lemmas 2.5 and 2.6 the system (1.2) has the exact and approximate solutions

$$y_n = \begin{cases} (1 + r)^{\frac{n}{2}} a, & n = 2, 4, 6, 8, \dots \\ (1 + r)^{\frac{n-1}{2}} b, & n = 1, 3, 5, 7, \dots \end{cases}$$

and

$$\psi_n = \begin{cases} (1 + r)^{\frac{n}{2}} a + \sum_{\lambda=0}^{\frac{n}{2}-1} (1 + r)^{\frac{n}{2}-1-\lambda} f_{2\lambda}, & n = 2, 4, 6, 8, \dots \\ (1 + r)^{\frac{n-1}{2}} b + \sum_{\lambda=0}^{\frac{n-1}{2}-1} (1 + r)^{\frac{n-1}{2}-1-\lambda} f_{2\lambda+1}, & n = 1, 3, 5, 7, \dots \end{cases}$$



Now if n is even, we have

$$\begin{aligned}
 \|y_n - \psi_n\| &= \left\| (1+r)^{\frac{n}{2}} a - (1+r)^{\frac{n}{2}} a - \sum_{\lambda=0}^{\frac{n}{2}-1} (1+r)^{\frac{n}{2}-1-\lambda} f_{2\lambda} \right\| \\
 &= \left\| \sum_{\lambda=0}^{\frac{n}{2}-1} (1+r)^{\frac{n}{2}-1-\lambda} f_{2\lambda} \right\| \\
 &\leq \varepsilon \sum_{\lambda=0}^{\frac{n}{2}-1} |1+r|^{\frac{n}{2}-1-\lambda} \\
 &\leq \varepsilon \sum_{\lambda=0}^{\infty} |1+r|^\lambda = \varepsilon(1+c^1+c^2+c^3+\dots) \\
 &= \varepsilon \frac{c}{1-c} \\
 &= \varepsilon \frac{1+r}{1-(1+r)} \\
 &= \varepsilon \frac{1+r}{-r} \\
 &= L\varepsilon.
 \end{aligned} \tag{3.1}$$

Now if n is odd, we have

$$\begin{aligned}
 \|y_n - \psi_n\| &= \left\| (1+r)^{\frac{n-1}{2}} b - (1+r)^{\frac{n-1}{2}} b - \sum_{\lambda=0}^{\frac{n-1}{2}-1} (1+r)^{\frac{n-1}{2}-1-\lambda} f_{2\lambda+1} \right\| \\
 &= \left\| \sum_{\lambda=0}^{\frac{n-1}{2}-1} (1+r)^{\frac{n-1}{2}-1-\lambda} f_{2\lambda+1} \right\| \\
 &\leq \varepsilon \sum_{\lambda=0}^{\frac{n-1}{2}-1} (1+r)^{\frac{n-1}{2}-1-\lambda} \\
 &\leq \varepsilon \sum_{\lambda=0}^{\infty} |1+r|^\lambda \\
 &= \varepsilon(1+c^1+c^2+c^3+\dots) \\
 &= \varepsilon \frac{c}{1-c} \\
 &= \varepsilon \frac{1+r}{1-(1+r)} \\
 &= \varepsilon \frac{1+r}{-r} \\
 &= L\varepsilon,
 \end{aligned} \tag{3.2}$$

where $L = \frac{1+r}{-r}$, hence the system is Hyers-Ulam stable. □

4. HYERS-ULAM STABILITY OF NON-AUTONOMOUS SYSTEMS

In this section we study the Hyers-Ulam stability of non-autonomous difference systems of first and second order. For the next two results we need the following assumptions:



$$\begin{aligned}\Lambda_3 : |r(n, \lambda + 1)| &\leq Me^{-\gamma(n-\lambda-1)}, \\ \Lambda_4 : \left| \prod_{\lambda=1}^j (r_{n-2\lambda}) \right| &< Me^{-\mu(j-1)}.\end{aligned}$$

Theorem 4.1. *If Λ_3 holds then, the system (1.3) is Hyer-Ulam stable.*

Proof. Since by Lemmas 2.5 and 2.6 the system (1.3) has the exact and approximate solutions

$$y_n = E(n, 0)\theta$$

and

$$\phi_n = E(n, 0)\theta + \sum_{\lambda=1}^n E(n, \lambda)f_{\lambda-1},$$

where $E(n, k)$ is the evolution family. Now consider

$$\begin{aligned}\|y_n - \phi_n\| &= \|r(n, 0)\theta - r(n, 0)\theta + \sum_{\lambda=0}^{n-2} r(n, \lambda + 1)f_{\lambda} + f_{n-1}\| \\ &= \left\| - \sum_{\lambda=0}^{n-2} r(n, \lambda + 1)f_{\lambda} + f_{n-1} \right\| = \left\| \sum_{\lambda=0}^{n-2} r(n, \lambda + 1)f_{\lambda} + f_{n-1} \right\| \\ &\leq \epsilon M \sum_{\lambda=0}^{n-2} e^{-\gamma(n-\lambda-1)} \\ &= \epsilon M (e^{-\gamma(n-1)} + e^{-\gamma(n-2)} + e^{-\gamma(n-3)} + \dots + 1) \\ &\leq \epsilon M (1 + e^{-\gamma} + e^{-2\gamma} + e^{-3\gamma} + e^{-4\gamma} + \dots) \\ &= \epsilon M \frac{1}{1 - e^{-\gamma}} = \epsilon M \frac{1}{1 - \frac{1}{e^\gamma}} \\ &= \epsilon M \frac{e^\gamma}{e^\gamma - 1} = \epsilon L,\end{aligned}\tag{4.1}$$

where $L = M \frac{e^\gamma}{e^\gamma - 1}$, hence the system (1.3) is Hyers-Ulam stable. \square

Theorem 4.2. *If Λ_4 holds then, the system (1.4) is Hyers-Ulam stable.*

Proof. Since by Lemmas 2.5 and 2.6 the system (1.4) has the exact and approximate solutions

$$y_n = \begin{cases} \prod_{\lambda=1}^{\frac{n}{2}} r_{n-2\lambda}\alpha, & n = 2, 4, 6, 8, \dots \\ \prod_{\lambda=1}^{\frac{n-1}{2}} r_{n-2\lambda}\beta, & n = 1, 3, 5, 7, \dots \end{cases}$$

and

$$\Psi_n = \begin{cases} \prod_{\lambda=1}^{\frac{n}{2}} (r_{n-2\lambda})\alpha + \sum_{j=1}^{\frac{n}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda})f_{n-2\lambda-2} + f_{n-2}, & n = 2, 4, 6, 8, \dots \\ \prod_{\lambda=1}^{\frac{n-1}{2}} (r_{n-2\lambda})\beta + \sum_{j=1}^{\frac{n-1}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda})f_{n-2\lambda-2} + f_{n-2}, & n = 1, 3, 5, 7, \dots \end{cases}$$



Now if n is even, we have

$$\begin{aligned}
 \|y_n - \Psi_n\| &= \left\| \prod_{\lambda=1}^{\frac{n}{2}} (r_{n-2\lambda})\alpha - \prod_{\lambda=1}^{\frac{n}{2}} (r_{n-2\lambda})\alpha - \sum_{j=1}^{\frac{n}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda})f_{n-2\lambda-2} + f_{n-2} \right\| \\
 &= \left\| \sum_{j=1}^{\frac{n}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda})f_{n-2\lambda-2} + f_{n-2} \right\| \leq \epsilon \left\| \sum_{j=1}^{\frac{n}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda}) \right\| \\
 &\leq \epsilon M \sum_{j=1}^{\frac{n}{2}-1} e^{-\mu(j-1)} = \epsilon M(1 + e^{-\mu} + e^{-2\mu} + e^{-3\mu} + \dots + e^{-(\frac{n-4}{2})\mu}) \\
 &\leq \epsilon M(1 + e^{-\mu} + e^{-2\mu} + e^{-3\mu} + \dots) = \epsilon M \frac{1}{1 - e^{-\mu}} = \epsilon M \frac{e^{\mu}}{e^{\mu} - 1} = \epsilon L.
 \end{aligned} \tag{4.2}$$

Now if n is odd, we have

$$\begin{aligned}
 \|y_n - \Psi_n\| &= \left\| \prod_{\lambda=1}^{\frac{n-1}{2}} (r_{n-2\lambda})\beta - \prod_{\lambda=1}^{\frac{n-1}{2}} (r_{n-2\lambda})\beta - \sum_{j=1}^{\frac{n-1}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda})f_{n-2\lambda-2} + f_{n-2} \right\| \\
 &= \left\| \sum_{j=1}^{\frac{n-1}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda})f_{n-2\lambda-2} + f_{n-2} \right\| \leq \epsilon \left\| \sum_{j=1}^{\frac{n-1}{2}-1} \prod_{\lambda=1}^j (r_{n-2\lambda}) \right\| \\
 &\leq \epsilon M \sum_{j=1}^{\frac{n-1}{2}-1} e^{-\mu(j-1)} = \epsilon M(1 + e^{-\mu} + e^{-2\mu} + e^{-3\mu} + \dots + e^{-(\frac{n-5}{2})\mu}) \\
 &\leq \epsilon M(1 + e^{-\mu} + e^{-2\mu} + e^{-3\mu} + \dots) = \epsilon M \frac{1}{1 - e^{-\mu}} = \epsilon M \frac{e^{\mu}}{e^{\mu} - 1} = \epsilon L,
 \end{aligned} \tag{4.3}$$

where $L = M \frac{e^{\mu}}{e^{\mu} - 1}$, hence the system is Hyers-Ulam stable. □

5. UNIFORM EXPONENTIAL STABILITY

In this section we study the uniform exponential stability of autonomous difference systems.

Theorem 5.1. *If Λ_1 holds then the system (1.1) is uniformly exponentially stable.*

Proof. Since system (1.1) has the solution

$$y_n = (1 + r)^n \gamma.$$

Consider

$$\|y_n\| = \|a^n \gamma\| \leq |\gamma| \|a\|^n \leq \xi e^{-\mu n} \tag{5.1}$$

where $a = 1 + r$ and $a^n = e^{\ln |a|^n} = e^{n \ln |a|} = e^{-\mu n}$, $0 < a < 1$, $\ln |a| = -\mu$, hence the system is uniformly exponentially stable. □

Theorem 5.2. *If Λ_1 holds then, the system (1.2) is uniformly exponentially stable.*

Proof. Since the system (1.2) has the solution:

$$y_n = \begin{cases} (1 + r)^{\frac{n}{2}} a, & n = 2, 4, 6, 8, \dots \\ (1 + r)^{\frac{n-1}{2}} b, & n = 1, 3, 5, 7, \dots \end{cases}$$

Now if n is even, we have

$$\begin{aligned}
 \|y_n\| &= \|(1 + r)^{\frac{n}{2}} a\| = \|\lambda^{\frac{n}{2}} a\| \leq |a| \|\lambda\|^{\frac{n}{2}} \leq |a| \|\lambda\|^n \\
 &\leq \Psi e^{-\mu n}.
 \end{aligned} \tag{5.2}$$



Now if n is odd, we have

$$\begin{aligned} \|y_n\| &= \|(1+r)^{\frac{n-1}{2}} b\| = \|\lambda^{\frac{n-1}{2}} b\| \leq |b| \|\lambda\|^{\frac{n-1}{2}} \leq |b| \|\lambda\|^n \\ &\leq \Psi e^{-\mu n}, \end{aligned} \quad (5.3)$$

where $\lambda = 1+r$ and $\lambda^n = e^{\ln|\lambda|^n} = e^{n \ln|\lambda|} = e^{-\mu n}$, $0 < \lambda < 1$ $\ln|\lambda| = -\mu$, hence the system is UES. \square

6. EXAMPLES

In this section we will present some examples which will elaborate the above results. For the autonomous and first order difference equation, we have the following example.

Example 6.1. Consider the autonomous difference equation:

$$\begin{cases} y_{n+1} = (1+r)y_n, & r = -0.5, n \geq 0 \\ y_0 = 0.3. \end{cases}$$

Then the following will be its solution:

$$y_n = (1+r)^n \gamma = (0.5)^n (0.3).$$

For approximate solution let $f_n = \varepsilon 0.6$ and consider the purterb system:

$$\begin{cases} \varphi_{n+1} = (1+r)\varphi_n + \varepsilon 0.6, & r = -0.5, n \geq 0 \\ \varphi_0 = 0.3. \end{cases}$$

Its solution is

$$\varphi_n = (0.5)^n (0.3) + \sum_{j=0}^{n-1} (0.5)^{n-1-j} (\varepsilon 0.6),$$

which is the approximate solution of the above system.

Now consider

$$\begin{aligned} \|y_n - \varphi_n\| &= \|(0.5)^n (0.3) - (0.5)^n (0.3) - \sum_{j=0}^{n-1} (0.5)^{n-1-j} (\varepsilon 0.6)\| \\ &\leq (\varepsilon 0.6) \sum_{j=0}^{n-1} (0.5)^{n-1-j} \\ &\leq (\varepsilon 0.6) \sum_{j=0}^{\infty} (0.5)^j = \frac{\varepsilon 0.6}{0.5} = \frac{6}{5} \varepsilon. \end{aligned}$$

Thus by Theorem 3.1 the above system is Hyer-Ulam stable.

Now for the uniform exponential stability we have, $\|y_n\| \leq 0.3e^{-\mu n}$, where $\mu = \ln 2$, since $e^{-\mu n} = (0.5)^n \Rightarrow \ln(e^{-\mu n}) = \ln(0.5)^n \Rightarrow -\mu n = n \ln(0.5) \Rightarrow -\mu = -\ln 2 \Rightarrow \mu = \ln 2$. Hence by Theorem 5.1 the system is also uniformly exponentially stable.

For second order autonomous system we have the following example.

Example 6.2. Consider the second order difference equation:

$$\begin{cases} y_{n+2} = \frac{1}{2}y_n, & n \geq 0 \\ y_0 = a \\ y_1 = b. \end{cases} \quad (6.1)$$



The solution is given as

$$y_n = \begin{cases} \left(\frac{1}{2}\right)^{\frac{n}{2}} a, & \text{when } n \text{ is even,} \\ \left(\frac{1}{2}\right)^{\frac{n-1}{2}} b, & \text{when } n \text{ is odd.} \end{cases}$$

Since the axiom λ_1 is satisfied so the system is Hyer-Ulam as well as uniformly exponentially stable.

The next example is for the case of non-autonomous system.

Example 6.3. Consider the non-autonomous difference equation:

$$\begin{cases} y_{n+1} = \frac{1}{2^n} y_n, & n \geq 0 \\ y_0 = 0.3. \end{cases}$$

Then the following will be its solution:

$$y_n = E(n, 0)\theta = \frac{1}{2^{n-1}} \frac{1}{2^{n-2}} \frac{1}{2^{n-3}} \cdots \frac{1}{2^0} \theta = \frac{1}{2^{\frac{n(n-1)}{2}}} \theta.$$

The approximate solution of the above system will be:

$$\phi_n = E(n, 0)\theta + \sum_{\lambda=1}^n E(n, \lambda) f_{\lambda-1} = \frac{1}{2^{\frac{n(n-1)}{2}}} \theta + \sum_{\lambda=1}^n \frac{1}{2^{\frac{n(n-1)-\lambda(\lambda-1)}{2}}} f_{\lambda-1},$$

where $\|f\| \leq \epsilon$.

Consider

$$\|y_n - \phi_n\| = \left\| \sum_{\lambda=1}^n \frac{1}{2^{\frac{n(n-1)-\lambda(\lambda-1)}{2}}} f_{\lambda-1} \right\| \leq \epsilon \sum_{\lambda=1}^n \frac{1}{2^\lambda} \leq \epsilon \sum_{\lambda=1}^{\infty} \frac{1}{2^\lambda} = 1 \cdot \epsilon,$$

hence by Theorem 4.1, the system is Hyers-Ulam stable.

For uniform exponential stability we have:

$$\|y_n\| = \left\| \frac{1}{2^{\frac{n(n-1)}{2}}} \theta \right\| \leq \|\theta\| \frac{1}{2^n},$$

which goes to zero exponentially hence by Theorem 5.1, the system is uniformly exponentially stable.

On the same way we can easily construct an example for non-autonomous second order difference system which we leave for the reader.

7. COCNLUSION

The study of difference equations is too much important and have a lot of applications in the different filed of sciences. The main purpose of this paper is to discussed the solutions, Hyer-Ulam and uniform exponential stabilities of autonomous and non-autonomous difference equations of first and second order. These models are not discussed yet before and therefore it can be fruitfull for other researcchers working in the field of difference equations and stability theory. Simple conditions were provided for each system through which one can easily check the stability of the proposed models. Some examples are also given to make sure the credibility of the main results.

ACKNOWLEDGMENTS

The authors would like to thank the Prince Sultan University for the support of this work through TAS LAB.



REFERENCES

- [1] S. Frassu and G. Viglialoro, *Boundedness for a fully parabolic Keller-Segel model with sublinear segregation and superlinear aggregation*, Acta Applicandae Mathematicae, 171(1) (2021), 1-20.
- [2] T. Li, N. Pintus, and G. Viglialoro, *Properties of solutions to porous medium problems with different sources and boundary conditions*, Zeitschrift für angewandte Mathematik und Physik, 70(3) (2019), 1-18.
- [3] T. Li and G. Viglialoro, *Boundedness for a non local reaction chemotaxis model even in the attraction-dominated regime*, Differential and Integral Equations, 34 (2021), 315-336.
- [4] T. Li and G. Viglialoro, *Analysis and explicit solvability of degenerate tensorial problems*, Boundary Value Problems, 2018(1) (2018), 1-13.
- [5] S. Brianzoni, C. Mammana, E. Michetti, and F. Zirilli, *A stochastic cobweb dynamical model*, Discrete Dynamics in Nature and Society, 2008 (2008).
- [6] J. Diblk, I. Dzhalladova, and M. Rikov, *Stabilization of companys income modeled by a system of discrete stochastic equations*, Advances in Difference Equations, 2014(1) (2014), 1-8.
- [7] S. M. Ulam, *A Collection of Mathematical Problems*, Inter science Publishers, 8 (1960).
- [8] D. H. Hyers, *On the stability of the linear functional equation*, Proceedings of the national academy of sciences of the United States of America, 27(4) (1941), 222.
- [9] T. Rassias, *On the stability of the linear mapping in Banach spaces*, Proceedings of the American mathematical society, 72(2) (1978), 297-300.
- [10] M. Obloza, *Connections between Hyers and Lyapunov stability of the ordinary differential equations*, (1997).
- [11] S. M. Jung, *Hyers-Ulam stability of the first order matrix difference equations*, Advances in Difference Equations, 2015(1) (2015), 1-13.
- [12] A. Khan, G. Rahmat, and A. Zada, *On uniform exponential stability and exact admissibility of discrete semigroups*, International Journal of Differential Equations, 2013 (2013).
- [13] Z. Gao, X. Yu, and J. Wang, *Exp-type Ulam-Hyers stability of fractional differential equations with positive constant coefficient*, Advances in Difference Equations, 2015(1) (2015), 1-10.
- [14] S. O. Shah, A. Zada, M. Muzammil, M. Tayyab, and R. Rizwan, *On the Bielecki-Ulam's type stability results of first order non-linear impulsive delay dynamic systems on time scales*, Qualitative Theory of Dynamical Systems, 19(3) (2020), 1-18.
- [15] A. R. Aruldass, D. Pachaiyappan, and C. Park, *HyersUlam stability of second-order differential equations using Mahgoub transform*, Advances in Difference Equations, 2021(1) (2021), 1-10.
- [16] A. Boutiara, S. Etemad, A. Hussain, and S. Rezapour, *The generalized Ulam-Hyer and UlamHyer stability and existence analysis of a coupled hybrid system of integro-differential IVPs involving ϕ -Caputo fractional operators*, Advances in Difference Equations, 2021(1) (2021), 1-21.
- [17] C. Buşe, A. Khan, G. Rahmat, and A. Tabassum, *A new estimation of the growth bound of a periodic evolution family on Banach spaces*, Journal of Function Spaces and Applications (Journal of function spaces), 2013 (2013), 6 pages.
- [18] S. Hussain, M. Sarwar, G. Rahmat, H. Aydi, and M. D. L Sen, *Mild solutions and controllability of fractional evolution inclusions of Clarkes subdifferential type with nonlocal conditions in Hilbert spaces*, Alexandria Engineering Journal, 80 (2023), 58-73.
- [19] S. Moonswan, G. Rahmat, A. Ullah, M. Y. Khan, Kamran, and K. Shah, *Hyers-Ulam stability, exponential stability and relative controllability of non-singular delay difference equations*, Complexity, 2022 (2022), Article ID. 8911621.
- [20] A. Lachouri, M. E. Samei, and A. Ardjouni, *Existence and stability analysis for a class of fractional pantograph q -difference equations with nonlocal boundary conditions*, Boundary Value Problems, 2023 (2023).
- [21] A. Boutiara, M. Benbachir, M. k. A. Kaabar et al. *Explicit iteration and unbounded solutions for fractional q -difference equations with boundary conditions on an infinite interval*, J Inequal Appl, 2022 (2022).
- [22] M. I. Ayari and S. T. M. Thabet, *Qualitative properties and approximate solutions of thermostat fractional dynamics system via a nonsingular kernel operator*, Arab Journal of Mathematical Sciences, (2023).



- [23] S. T. M. Thabet, M. V. Cortez, I. Kedim, M. E. Samei, and M. I. Ayari, *Solvability of ρ -Hilfer fractional snap dynamic system on unbounded domains*, *Fractals and Fractional*, (2023).
- [24] Y. Almalki, G. Rahmat, A. Ullah, F. Shehryar, M. Numan, and M. U. Ali, *Generalized β -Hyers-Ulam-Rassias stability of impulsive difference equations*, *Computational Intelligence and Neurosciences, 2022* (2022), Article ID. 9462424.
- [25] O. Tunç and C. Tunç, *On Ulam stabilities of iterative Fredholm and Volterra integral equations with multiple time-varying delays*, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.*, *118*(3) (2024), 20 pp.
- [26] O. Tunç and C. Tunç, *Ulam stabilities of nonlinear iterative integro-differential equations*, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.*, *117*(3) (2023).
- [27] O. Tunç and C. Tunç, *On Ulam stabilities of delay Hammerstein integral equation*, *Symmetry*, *2023*(15), 1736.
- [28] O. Tunç, D. R. Sahu, and C. Tunç, *On the Ulam type stabilities of a general iterative integro-differential equation including a variable delay*, *J. Nonlinear Convex Anal.*, *25*(2) (2024), 399417.
- [29] O. Tunç, C. Tunç, and J. -C. Yao, *New results on Ulam stabilities of nonlinear integral equations*, *Mathematics*, *2024*(12) (2024), 682.
- [30] O. Tunç, C. Tunç, G. Petruşel, and J. -C. Yao, *On the Ulam stabilities of nonlinear integral equations and integro-differential equations*, *Math. Meth. Appl. Sci.*, *47*(6) (2024), 115.

Uncorrected Proof

