



Dynamic insights into gaseous diffusion: analytical soliton and wave solutions via Chaffee–Infante equation in homogeneous media

Samah Mabrouk¹, Mahy Mahdy¹, Rasha Saleh¹, and Ahmed Rashed^{1,2,*}

¹Department of Physics and Engineering Mathematics, Faculty of Engineering, Zagazig University, Egypt.

²Basic Science department, Faculty of Engineering, Delta University for Science and Technology, 11152, Gamasa, Egypt.

Abstract

Gaseous diffusion (GD) has been used in various fields, including electromagnetic wave fields, high-energy physics, fluid dynamics, coastal engineering, ion-acoustic waves in plasma physics, and optical fibers. GD involves random molecular movement from areas of high partial pressure to areas of low partial pressure. Researchers have developed models to describe this phenomenon, among these models is the $(2 + 1)$ -dimensional Chaffee–Infante (CI)-equation. This research explores analytical soliton and wave solutions of Gaseous diffusion through a homogeneous medium, considering two analytical methods, the Riccati equation and F-expansion methods. Thirty-seven different solutions have been identified and some of these solutions have been illustrated graphically. The figures show a range of bright, dark, singular, singular-periodic, and kink-type soliton wave solutions.

Keywords. $(2 + 1)$ -dimensional Chaffee–Infante, Gaseous diffusion (GD), Riccati equation method, F-expansion method.

2010 Mathematics Subject Classification. 58J47, 35C05.

1. INTRODUCTION

One of the most significant natural phenomena, widely used in numerous fields such as physics, biology, and chemistry is gaseous diffusion. Gaseous diffusion is the movement of molecules under a concentration gradient. GD has produced most of the enriched uranium in the world [1]. GD in materials science is widespread, for example, in processes such as sintering, corrosion, steel hardening, and semiconductor manufacturing. The diffusion of gas in a homogeneous medium involves studying a particle with a homogeneous temperature under the influence of an external force. Scientists have developed models to describe this phenomenon, among which is the $(2 + 1)$ -dimensional Chaffee–Infante (CI) equation [2]. CI equation can depict the physical phenomena of particle diffusion, which has been extensively used in electromagnetic wave fields, fluid dynamics, high-energy physics, coastal engineering, fluid mechanics, and ion-acoustic waves in plasma physics, optical fibers, and other fields [1]. CI equation can depict the physical phenomena of a particle diffusion, which has been extensively used in electromagnetic wave fields, fluid dynamics, high-energy physics, coastal engineering, fluid mechanics, and ion-acoustic waves in plasma physics, optical fibers, and other fields [1]. There are many techniques that can be used to solve a wide range of higher-dimensional nonlinear equations in the applied sciences and mathematical physics such as Lie and symmetry analysis [2–14], the inverse scattering transformation method [15], the Darboux transformation method [16, 17], generalized exponential rational function (GERF) technique [1, 18, 19], the Riccati equation method [20, 21], the (G/G) expansion [23–25], the tanh-coth (TC) method [26, 27], F-expansion method [28, 29], the Backlund transformation method [30, 31], and extra methods have been developed to get the exact solutions of nonlinear equations. Recently CI equation has been widely studied by many researchers. Khater and Ghanbari utilized five methods to obtain the solitary wave solutions for the CI equation: the $\exp(-\phi)$ -expansion method, the extended (G'/G) -expansion method, the extended simplest equation method, the extended tanh expansion method, and the modified Khater method [32]. Moreover,

Received: 18 July 2024; Accepted: 04 November 2024.

* Corresponding author. Email: ahmed.s.rashed@gmail.com.

the differential quadrature method (DQM) is also an effective technique [33]. Also, many of these techniques are very effective in the case of fractional calculus. Other techniques can be found here [34–36]. In [1], the GERF method is used to produce a great number of analytical soliton solutions. According to Akbar et al. [37], the first integral method has been employed to find the analytic solutions of the CI equation. In [38], the authors used the modified Khatter method to seek the CI equation with numerous new closed-form solutions. The solitary wave solution is found by the extended sinh-Gordon expansion technique [39]. There are various other techniques for the construction of different kinds of solutions. In this article, we will apply two distinct methods, namely the Riccati equation and the F-expansion method, to find the analytic solutions of the CI-equation, which has the form:

$$\psi_{xt} + (-\psi_{xx} + \alpha\psi^3 - \alpha\psi)_x + \sigma\psi_{yy} = 0, \quad (1.1)$$

where, α is the diffusion coefficient and σ is the degradation coefficient.

The paper is arranged as follows. Section 2 introduces the Riccati equation method with application to the $(2 + 1)$ -dimensional Chaffee–Infante equation. In section 3, the soliton wave solutions of the CI- equation are argued via F-expansion method. The paper ends with conclusions in section 4.

2. THE RICCATI EQUATION METHOD FOR THE $(2 + 1)$ -DIMENSIONAL CI-EQUATION

In this section, the Riccati equation method is discussed and applied to find the soliton solutions of the CI- equation. The method can be summarized in the following steps:

Step 1: Reduce the order of the partial differential equation via wave transformation. Consider

$$\psi(x, y, t) = \psi(\tau), \quad \tau = \mu x + \lambda y - \gamma t. \quad (2.1)$$

Substituting Eq. (2.1) into (1.1) reduces it to:

$$-\mu^3\psi''' - \mu\alpha\psi' + 3\mu\alpha\psi^2\psi' - \mu\gamma\psi'' + \sigma\lambda^2\psi'' = (-\mu^3\psi'' - \mu\alpha\psi + \mu\alpha\psi^3)' + (\sigma\lambda^2 - \mu\gamma)\psi'' = 0. \quad (2.2)$$

Which can be simplified after integration to:

$$-\mu^3\psi'' + (\sigma\lambda^2 - \mu\gamma)\psi' - \mu\alpha\psi + \mu\alpha\psi^3 = 0. \quad (2.3)$$

Step 2: Assume that the solution of the reduced equation is in a series form:

$$\psi(\tau) = \sum_{i=0}^N A_i \phi^i(\tau) \quad (2.4)$$

where A_i are real constants will be determined, N is a positive integer, which result from the balancing principle with higher order non-linear and linear terms in Eq. (2.3), by balancing ψ'' and ψ^3 gives $N=1$, and Eq. (2.4) is written as:

$$\psi(\tau) = A_0 + A_1\phi, \quad (2.5)$$

as $\phi(\tau)$ satisfies the following Riccati equation:

$$\phi'(\tau) = a\phi^2(\tau) + b\phi + c \quad (2.6)$$

where $a \neq 0$, b and c will be determined later. The solutions of the Riccati equation can be written as follows:

Case (1): $\Omega > 0$,

$$\phi(\xi) = -\frac{b}{2a} - \frac{\sqrt{\Omega}}{2a} \tanh\left(\frac{\sqrt{\Omega}}{2}\xi + \xi_0\right), \quad (2.7)$$

$$\phi(\xi) = -\frac{b}{2a} - \frac{\sqrt{\Omega}}{2a} \coth\left(\frac{\sqrt{\Omega}}{2}\xi + \xi_0\right). \quad (2.8)$$

Case (2): $\Omega < 0$,

$$\phi(\xi) = -\frac{b}{2a} - \frac{\sqrt{-\Omega}}{2a} \tan\left(\frac{\sqrt{-\Omega}}{2}\xi + \xi_0\right), \quad (2.9)$$



$$\phi(\xi) = -\frac{b}{2a} - \frac{\sqrt{-\Omega}}{2a} \cot\left(\frac{\sqrt{-\Omega}}{2}\xi + \xi_0\right). \quad (2.10)$$

Case (3): $\Omega = 0$,

$$\phi(\xi) = -\frac{b}{2a} - \frac{1}{a\xi + \xi_0}, \quad (2.11)$$

where $\Omega = b^2 - 4ac$, and ξ_0 is the integration constant. **Step 3:** Inserting Eq. (2.5) together with Eq. (2.6) into Eq. (2.3), collecting all coefficients of each power of f^i , $0 \leq i \leq N$ in the resulting equation where these coefficients must vanish. This gives a system of algebraic equations involving the parameters A_i , ($i = 1, 2, 3$), a , b , μ , λ , γ , and c

$$\begin{cases} \psi = A_0 + A_1\varphi, \\ \psi' = A_1\varphi' = aA_1\varphi^2 + bA_1\varphi + cA_1, \\ \psi'' = A_1\varphi'' = 2a^2A_1\varphi^3 + 3abA_1\varphi^2 + (b^2A_1 + 2acA_1)\varphi + cbA, \\ \psi^3 = A_1^3\varphi^3 + 3A_0A_1^2\varphi^2 + 3A_0^2A_1\varphi + A_0^3. \end{cases} \quad (2.12)$$

Substituting (2.12) into (2.3), gathering coefficients of each φ^i power, and setting the sum to zero yields the subsequent algebraic equations:

$$\text{Coefficient of } \varphi^3 = -2\mu^3a^2A_1 + \mu\alpha A_1^3, \quad (2.13)$$

$$\text{Coefficient of } \varphi^2 = -3\mu^3abA_1 + aA_1\sigma\lambda^2 - kmaA_1 + 3\mu\alpha A_0A_1^2, \quad (2.14)$$

$$\text{Coefficient of } \varphi = -k^3b^2A_1 - 2\mu^3acA_1 + bA_1\sigma\lambda^2 - bA_1\mu\gamma + 3\mu\alpha A_0^2A_1 - \mu\alpha A_1, \quad (2.15)$$

$$\text{Coefficient of } \varphi^0 = -\mu^3cbA + cA_1\sigma\lambda^2 - cA_1\mu\gamma + \mu\alpha A_0^3 - \mu\alpha A_0. \quad (2.16)$$

Solving the algebraic system (2.13)-(2.16) using the Maple package confers six different groups of solutions of the system, which result in twelve different solutions of the CI Equation (1.1) as:

Group1:

$$A_0 = \frac{b\mu}{\sqrt{2\alpha}}, \quad A_1 = \sqrt{\frac{2}{\alpha}}\mu a, \quad c = \frac{b^2\mu^2 - 2\alpha}{4a\mu^2}, \quad \gamma = \frac{\sigma\lambda^2}{\mu}, \quad \text{and} \quad \Omega = \frac{2\alpha}{\mu^2}. \quad (2.17)$$

Case1: $\Omega > 0$

$$\psi_1(\tau) = -\tanh\left(\frac{\sqrt{2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.18)$$

$$\psi_2(\tau) = -\coth\left(\frac{\sqrt{2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.19)$$

The kink wave ψ_1 is presented for $a = 1$, $b = 3$, $c = 1.25$, $\alpha = 2$, $t = 1$, and $\gamma = \lambda = \mu = 1$ at Figure 1, while the singular-type soliton solution (2.19) at $a = 1$, $b = 3$, $c = 1.25$, $\alpha = 2$, $t = 1$, and $\gamma = \lambda = \mu = 1$ at Figure 2.

Case (2): $\Omega < 0$,

$$\psi_3(\tau) = i \tan\left(\frac{\sqrt{-2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.20)$$

$$\psi_4(\tau) = -i \cot\left(\frac{\sqrt{-2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.21)$$

Figure 3 indicates the double soliton solution (2.20) at $a = 1$, $b = 3$, $c = 1.25$, $\alpha = -10$, and $-\gamma = \lambda = \mu = 5$.

Group2:






FIGURE 1. Kink wave ψ_1 at $a = 1$, $b = 3$, $c = 1.25$, $\alpha = 2$, $t = 1$, and $\gamma = \lambda = \mu = 1$.




FIGURE 2. singular soliton solution ψ_2 at $a = 1$, $b = 3$, $c = 1.25$, $\alpha = 2$, $t = 1$, and $\gamma = \lambda = \mu = 1$.




FIGURE 3. The double soliton solution ψ_3 at $a = 1$, $b = 3$, $c = 1.25$, $\alpha = 2$, $t = 1$, and $\gamma = \lambda = \mu = 1$.




FIGURE 4. solitary wave solution ψ_6 at $a = 1$, $b = 3$, $c = 2$, $\alpha = 2$, $t = 1$, and $\gamma = 4$, $\lambda = \mu = 1$.


$$A_0 = \frac{\sqrt{2}b\mu + \sqrt{\alpha}}{2\sqrt{\alpha}}, \quad A_1 = \sqrt{\frac{2}{\alpha}}\mu a, \quad c = \frac{2b^2\mu^2 - \alpha}{8a\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 + \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad \text{and} \quad \Omega = \frac{\alpha}{2\mu^2}. \quad (2.22)$$

Case1: $\Omega > 0$

$$\psi_5(\tau) = \frac{1}{2} - \frac{1}{2} \tanh \left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu} (\mu x + \lambda y - \gamma t) \right), \quad (2.23)$$

$$\psi_6(\tau) = \frac{1}{2} - \frac{1}{2} \coth \left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu} (\mu x + \lambda y - \gamma t) \right). \quad (2.24)$$



FIGURE 5. ψ_8 at $a = 1$, $b = 3$, $c = 2$, $\alpha = 2$, $t = 1$, and $\gamma = 4$, $\lambda = \mu = 1$.

The solitary wave solution ψ_6 at $a = 1$, $b = 3$, $c = 2$, $\alpha = 2$, $t = 1$ and $\gamma = 4$, $\lambda = \mu = 1$. is shown in Figure 4.

Case (2): $\Omega < 0$,

$$\psi_7(\tau) = \frac{1}{2} + \frac{1}{2}i \tan \left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t) \right), \quad (2.25)$$

$$\psi_8(\tau) = \frac{1}{2} - \frac{1}{2}i \cot \left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t) \right). \quad (2.26)$$

The solution, ψ_8 , at $a = 1$, $b = 3$, $c = 2$, $\alpha = 2$, $t = 1$, and $\gamma = 4$, $\lambda = \mu = 1$ is depicted in Figure 5.

Group3:

$$A_0 = \frac{-b\mu}{\sqrt{2\alpha}}, \quad A_1 = -\sqrt{\frac{2}{\alpha}}\mu a, \quad c = \frac{b^2\mu^2 - 2\alpha}{4a\mu^2}, \quad \gamma = \frac{\sigma\lambda^2}{\mu}, \quad \text{and} \quad \Omega = \frac{2\alpha}{\mu^2}. \quad (2.27)$$

Case1: $\Omega > 0$

$$\psi_9(\tau) = \tanh \left(\frac{\sqrt{2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t) \right), \quad (2.28)$$

$$\psi_{10}(\tau) = \coth \left(\frac{\sqrt{2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t) \right). \quad (2.29)$$

Case (2): $\Omega < 0$,

$$\psi_{11}(\tau) = -i \tan \left(\frac{\sqrt{-2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t) \right), \quad (2.30)$$

$$\psi_{12}(\tau) = i \cot \left(\frac{\sqrt{-2\alpha}}{2\mu}(\mu x + \lambda y - \gamma t) \right). \quad (2.31)$$

Group4:

$$A_0 = \frac{\sqrt{2}b\mu - \sqrt{\alpha}}{2\sqrt{\alpha}}, \quad A_1 = \sqrt{\frac{2}{\alpha}}\mu a, \quad c = \frac{2b^2\mu^2 - \alpha}{8a\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 - \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad \Omega = \frac{\alpha}{2\mu^2}. \quad (2.32)$$



Case1: $\Omega > 0$

$$\psi_{13}(\tau) = -\frac{1}{2} - \frac{1}{2} \tanh\left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.33)$$

$$\psi_{14}(\tau) = -\frac{1}{2} - \frac{1}{2} \coth\left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right) \quad (2.34)$$

Case (2): $\Omega < 0$,

$$\psi_{15}(\tau) = -\frac{1}{2} + \frac{1}{2}i \tan\left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.35)$$

$$\psi_{16}(\tau) = -\frac{1}{2} - \frac{1}{2}i \cot\left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.36)$$

Group 5:

$$A_0 = \frac{-\sqrt{2}b\mu - \sqrt{\alpha}}{2\sqrt{\alpha}}, \quad A_1 = -\sqrt{\frac{2}{\alpha}}\mu a, \quad c = \frac{2b^2\mu^2 - \alpha}{8a\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 + \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad \text{and} \quad \Omega = \frac{\alpha}{2\mu^2}. \quad (2.37)$$

Case1: $\Omega > 0$

$$\psi_{17}(\tau) = -\frac{1}{2} + \frac{1}{2} \tanh\left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.38)$$

$$\psi_{18}(\tau) = -\frac{1}{2} + \frac{1}{2} \coth\left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.39)$$

Case (2): $\Omega < 0$,

$$\psi_{19}(\tau) = -\frac{1}{2} - \frac{1}{2}i \tan\left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.40)$$

$$\psi_{20}(\tau) = -\frac{1}{2} + \frac{1}{2}i \cot\left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.41)$$

Group 6:

$$A_0 = \frac{-\sqrt{2}b\mu + \sqrt{\alpha}}{2\sqrt{\alpha}}, \quad A_1 = -\sqrt{\frac{2}{\alpha}}\mu a, \quad c = \frac{2b^2\mu^2 - \alpha}{8a\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 - \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad \Omega = \frac{\alpha}{2\mu^2}. \quad (2.42)$$

Case1: $\Omega > 0$

$$\psi_{21}(\tau) = \frac{1}{2} + \frac{1}{2} \tanh\left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad (2.43)$$

$$\psi_{22}(\tau) = \frac{1}{2} + \frac{1}{2} \coth\left(\frac{\sqrt{\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.44)$$

Case (2): $\Omega < 0$,

$$\psi_{23}(\tau) = \frac{1}{2} - \frac{1}{2}i \tan\left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right), \quad \psi_{24}(\tau) = \frac{1}{2} + \frac{1}{2}i \cot\left(\frac{\sqrt{-\frac{\alpha}{2}}}{2\mu}(\mu x + \lambda y - \gamma t)\right). \quad (2.45)$$



Case (3): $\Omega = 0$

$$\psi_{25}(\tau) = -\frac{1}{a(\mu x + \lambda y - \gamma t)}. \quad (2.46)$$

3. F-EXPANSION METHOD FOR THE $(2 + 1)$ - DIMENSIONAL CI- EQUATION

F-expansion method [21, 40] mainly starts from the reduced Equation (2.3), where the solution of the ODE is assumed to be:

$$\psi(\tau) = \sum_{i=0}^N s_i F^i(\tau), \quad (3.1)$$

where, $N = 1$. This value is obtained from the balancing between the higher-order nonlinear and linear terms in Eq. (2.3) and leads to:

$$\begin{cases} \psi = s_0 + s_1 F, \\ \psi' = s_1 F' = s_1 (pF^4 + QF^2 + R)^{0.5}, \\ \psi'' = s_1 F'' = 2s_1 pF^3 + s_1 QF, \\ \psi^3 = s_0^3 + 3s_0^2 s_1 F + 3s_0 s_1^2 F^2 + s_1^3 F^3. \end{cases} \quad (3.2)$$

s_i are real constants and $F(\tau)$ satisfies the following first order ODE:

$$F'(\xi) = (PF^4(\xi) + QF^2 + R)^{0.5}, \quad (3.3)$$

as $P \neq 0$, Q and R are real constants. They will be determined later. The solutions of Eq. (3.3) are illustrated in terms of Jacobian elliptic functions, for more details see [41]. Inserting Eq. (3.1) together with Eq. (3.3) into Eq. (2.3), collecting all coefficients of each power of F^i , $i = 0, 1, \dots, N$ to zero, results in the following system of algebraic equations:

$$\text{Coefficient of } F^6: 4p^2 s_1^2 \mu^6 - 4\alpha p s_1^4 \mu^4 + \alpha^2 \mu^2 s_1^6 = 0, \quad (3.4)$$

$$\text{Coefficient of } F^5: -12\alpha p s_0 s_1^3 \mu^4 + 6s_0 \alpha^2 \mu^2 s_1^5 = 0, \quad (3.5)$$

$$\begin{aligned} \text{Coefficient of } F^4: 4Qps_1^2 \mu^6 - 12\alpha p s_0^2 s_1^2 \mu^4 - 2Q\alpha s_1^4 \mu^4 + 15\alpha^2 s_0^2 \mu^2 s_1^4 + 4\alpha p s_1^2 \mu^4 \\ - 2\alpha^2 \mu^2 s_1^4 - p s_1^2 \sigma^2 \lambda^4 + 2p\mu\gamma\sigma s_1^2 \lambda^2 - p s_1^2 \mu^2 \gamma^2 = 0, \end{aligned} \quad (3.6)$$

$$\text{Coefficient of } F^3: -4\alpha p s_1 s_0^3 \mu^4 - 6Q\alpha s_0 s_1^3 \mu^4 + 20\alpha^2 \mu^2 s_0^3 s_1^3 + 4p s_0 s_1 \alpha \mu^4 - 8s_0 \alpha^2 \mu^2 s_1^3 = 0, \quad (3.7)$$

$$\begin{aligned} \text{Coefficient of } F^3: -4\alpha p s_1 s_0^3 \mu^4 - 6Q\alpha s_0 s_1^3 \mu^4 + 20\alpha^2 \mu^2 s_0^3 s_1^3 + 4p s_0 s_1 \alpha \mu^4 - 8s_0 \alpha^2 \mu^2 s_1^3 \\ - Qs_1^2 \sigma^2 \lambda^4 + 2Q\mu\gamma\sigma s_1^2 \lambda^2 - Qs_1^2 \mu^2 \gamma^2 = 0, \end{aligned} \quad (3.8)$$

$$\text{Coefficient of } F^1: -2Q\alpha s_1 s_0^3 \mu^4 + 6s_1 \alpha^2 \mu^2 s_0^5 + 2\alpha Q s_0 s_1 \mu^4 - 8s_1 \alpha^2 \mu^2 s_0^3 + 2s_0 s_1 \alpha^2 \mu^2 = 0, \quad (3.9)$$

$$\text{Coefficient of } F^0: \alpha^2 \mu^2 s_0^6 - 2\alpha^2 \mu^2 s_0^4 + \alpha^2 \mu^2 s_0^2 - R s_1^2 \sigma^2 \lambda^4 + 2R\mu\gamma\sigma s_1^2 \lambda^2 - R s_1^2 \mu^2 \gamma^2 = 0, \quad (3.10)$$

Solving the algebraic system (3.4)-(3.10) using the MAPLE package confers five different groups of solutions of the system which result in numerous different solutions of the CI Equation (1.1) as:

Group 1:

$$s_0 = 0, \quad s_1 = s_1, \quad p = \frac{s_1^2 \alpha}{2\mu^2}, \quad Q = \frac{-\alpha}{\mu^2}, \quad \gamma = \frac{\sigma \lambda^2}{\mu}, \quad (3.11)$$

Case 1: If $P = m^2$, $Q = -1 - m^2$, $m = 1$,

A kink-type solution is obtained:

$$\psi_{26}(\tau) = (s_1 \tanh(\mu x + \lambda y - \gamma t)). \quad (3.12)$$

This solution is plotted in Figure 6 at:

$$p = 1, \quad Q = -2, \quad R = 1, \quad \alpha = 2, \quad t = 1, \quad \gamma = \lambda = \mu = 1, \quad \text{and } s_1 = 1. \quad (3.13)$$






FIGURE 6. Kink wave ψ_{13} at $p = 1$, $Q = -2$, $R = 1$, $\alpha = 2$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = 1$.




FIGURE 7. Bright kink soliton solution ψ_{14} at $p = -1$, $Q = 1$, $R = 1$, $\alpha = -1$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = \sqrt{2}$.




FIGURE 8. Singular solution ψ_{15} at $p = 1$, $Q = -2$, $R = 1$, $\alpha = 2$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = 1$.

Case 2: If $P = -m^2$, $Q = 2m^2 - 1$, $m = 1$,
Bright soliton solutions is obtained:

$$\psi_{27}(\tau) = s_1 (\text{sech}(\mu x + \lambda y - \gamma t)) \quad (3.14)$$

This solution is illustrated in Figure 7 at $p = -1$, $Q = 1$, $R = 1$, $\alpha = -1$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = \sqrt{2}$.

Case 3: If $P = 1$, $Q = -1 - m^2$,

The following solutions are obtained:

a- Singular solution with $m = 1$

$$\psi_{28}(\tau) = s_1 (\coth(\mu x + \lambda y - \gamma t)), \quad (3.15)$$

which is graphed at $p = 1$, $Q = -2$, $R = 1$, $\alpha = 2$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = 1$, in Figure 8.

b-Singular periodic solution with $m = 0$

$$\psi_{29}(\tau) = s_1 (\csc(\mu x + \lambda y - \gamma t)). \quad (3.16)$$

This solution is depicted in Figure 9 at $p = 1$, $Q = -1$, $R = 1$, $\alpha = 1$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = \sqrt{2}$.

Case 4: If $P = 1 - m^2$, $Q = 2m^2 - 1$, $m = 0$






FIGURE 9. Singular periodic solution ψ_{16} at $p = 1$, $Q = -1$, $R = 1$, $\alpha = 1$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = \sqrt{2}$.




FIGURE 10. Singular periodic solution ψ_{17} at $p = 1$, $Q = -1$, $R = 1$, $\alpha = 1$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = \sqrt{2}$.

A singular periodic solution is obtained as:

$$\psi_{30}(\tau) = s_1 (\sec(\mu x + \lambda y - \gamma t)). \quad (3.17)$$

This solution is illustrated in Figure 10 for $p = 1$, $Q = -1$, $R = 1$, $\alpha = 1$, $t = 1$, $\gamma = \lambda = \mu = 1$, and $s_1 = \sqrt{2}$.

Group 2:

$$s_0 = \frac{1}{2}, \quad s_1 = s_1, \quad p = \frac{s_1^2 \alpha}{2\mu^2}, \quad Q = \frac{-\alpha}{4\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 + \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad R = \frac{\alpha}{32\mu^2 s_1^2}. \quad (3.18)$$

Case 1: If $P = m^2$, $Q = -1 - m^2$, $m = 1$.

A kink-type soliton solution is obtained:

$$\psi_{31}(\tau) = \frac{1}{2} + (s_1 \tanh(\mu x + \lambda y - \gamma t)). \quad (3.19)$$

Case 2: If $P = -m^2$, $Q = 2m^2 - 1$, $m = 1$.

A Bright and dark soliton solution is obtained:

$$\psi_{32}(\tau) = \frac{1}{2} + s_1 (\operatorname{sech}(\mu x + \lambda y - \gamma t)). \quad (3.20)$$

Case 3: If $P = 1$, $Q = -1 - m^2$.

The following solutions are obtained:

a-Singular solution with $m = 1$

$$\psi_{33}(\tau) = \frac{1}{2} + s_1 (\coth(\mu x + \lambda y - \gamma t)), \quad (3.21)$$

b-Singular periodic solution with $m = 0$.

$$\psi_{34}(\tau) = \frac{1}{2} + s_1 (\csc(\mu x + \lambda y - \gamma t)). \quad (3.22)$$

Case 4: If $P = 1 - m^2$, $Q = 2m^2 - 1$, $m = 0$.

A singular periodic solution is obtained:

$$\psi_{35}(\tau) = \frac{1}{2} + s_1 (\sec(\mu x + \lambda y - \gamma t)). \quad (3.23)$$



Group 3:

$$s_0 = \frac{-1}{2}, \quad s_1 = s_1, \quad p = \frac{s_1^2 \alpha}{2\mu^2}, \quad Q = \frac{-\alpha}{4\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 + \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad R = \frac{\alpha}{32\mu^2 s_1^2}. \quad (3.24)$$

Case 1: If $P = m^2$, $Q = -1 - m^2$, $m = 1$.

A kink-type soliton solution is obtained:

$$\psi_{36}(\tau) = -\frac{1}{2} + (s_1 \tanh(\mu x + \lambda y - \gamma t)). \quad (3.25)$$

Case 2: If $P = -m^2$, $Q = 2m^2 - 1$, $m = 1$.

Bright and dark soliton solutions is obtained:

$$\psi_{37}(\tau) = -\frac{1}{2} + s_1 (\operatorname{sech}(\mu x + \lambda y - \gamma t)). \quad (3.26)$$

Case 3: If $P = 1$, $Q = -1 - m^2$.

The following solutions are created:

a-Singular solution with $m = 1$

$$\psi_{38}(\tau) = -\frac{1}{2} + s_1 (\coth(\mu x + \lambda y - \gamma t)), \quad (3.27)$$

b-Singular periodic solution with $m = 0$

$$\psi_{39}(\tau) = -\frac{1}{2} + s_1 (\csc(\mu x + \lambda y - \gamma t)). \quad (3.28)$$

Case 4: If $P = 1 - m^2$, $Q = 2m^2 - 1$, $m = 0$.

A singular periodic solution is created:

$$\psi_{40}(\tau) = -\frac{1}{2} + s_1 (\sec(\mu x + \lambda y - \gamma t)). \quad (3.29)$$

Group 4:

$$s_0 = \frac{1}{2}, \quad s_1 = s_1, \quad p = \frac{s_1^2 \alpha}{2\mu^2}, \quad Q = \frac{-\alpha}{4\mu^2}, \quad \gamma = \frac{\sigma\lambda^2 - \frac{3}{\sqrt{2}}\sqrt{\alpha}\mu^2}{\mu}, \quad R = \frac{\alpha}{32\mu^2 s_1^2}, \quad (3.30)$$

Case 1: If $P = m^2$, $Q = -1 - m^2$, $m = 1$.

A kink-type soliton solution is obtained:

$$\psi_{41}(\tau) = \frac{1}{2} + (s_1 \tanh(\mu x + \lambda y - \gamma t)). \quad (3.31)$$

Case 2: If $P = -m^2$, $Q = 2m^2 - 1$, $m = 1$

Bright and dark soliton solution is given:

$$\psi_{42}(\tau) = \frac{1}{2} + s_1 (\operatorname{sech}(\mu x + \lambda y - \gamma t)). \quad (3.32)$$

Case 3: If $P = 1$, $Q = -1 - m^2$.

The following solutions are: *a-Singular solution with $m = 1$*

$$\psi_{43}(\tau) = \frac{1}{2} + s_1 (\coth(\mu x + \lambda y - \gamma t)), \quad (3.33)$$

b-Singular periodic solution with $m = 0$.

$$\psi_{44}(\tau) = \frac{1}{2} + s_1 (\csc(\mu x + \lambda y - \gamma t)). \quad (3.34)$$



Case 4: If $P = 1 - m^2, Q = 2m^2 - 1, m = 0$.

A singular periodic solution is given in the following form:

$$\psi_{45}(\tau) = \frac{1}{2} + s_1 (\sec(\mu x + \lambda y - \gamma t)). \quad (3.35)$$

Group 5:

$$s_0 = \frac{-1}{2}, \quad s_1 = s_1, \quad p = \frac{s_1^2 \alpha}{2\mu^2}, \quad Q = \frac{-\alpha}{4\mu^2}, \quad \gamma = \frac{\sigma \lambda^2 - \frac{3}{\sqrt{2}} \sqrt{\alpha} \mu^2}{\mu}, \quad R = \frac{\alpha}{32\mu^2 s_1^2}, \quad (3.36)$$

Case 1: If $P = m^2, Q = -1 - m^2, m = 1$.

A kink-type soliton solution is created:

$$\psi_{46}(\tau) = -\frac{1}{2} + (s_1 \tanh(\mu x + \lambda y - \gamma t)). \quad (3.37)$$

Case 2: If $P = -m^2, Q = 2m^2 - 1, m = 1$

Bright and dark soliton solution is given:

$$\psi_{47}(\tau) = -\frac{1}{2} + s_1 (\operatorname{sech}(\mu x + \lambda y - \gamma t)). \quad (3.38)$$

Case 3: If $P = 1, Q = -1 - m^2$.

The following solutions are created:

a-Singular solution with $m = 1$.

$$\psi_{48}(\tau) = -\frac{1}{2} + s_1 (\coth(\mu x + \lambda y - \gamma t)), \quad (3.39)$$

b-Singular periodic solution with $m = 0$.

$$\psi_{49}(\tau) = -\frac{1}{2} + s_1 (\csc(\mu x + \lambda y - \gamma t)). \quad (3.40)$$

Case 4: If $P = 1 - m^2, Q = 2m^2 - 1, m = 0$.

A singular periodic solution is obtained:

$$\psi_{50}(\tau) = -\frac{1}{2} + s_1 (\sec(\mu x + \lambda y - \gamma t)). \quad (3.41)$$

4. CONCLUSIONS

In this article, the dynamical behavior of gas diffusion was examined by investigating the $(2 + 1)$ -dimensional Chaffee–Infante equation. Firstly, the Riccati equation method is applied, revealing six groups of soliton solutions. The F-expansion method is then applied, which confers five different groups of solutions, resulting in twenty-five different solutions. The obtained solutions vary between kink-type soliton, singular soliton, singular periodic dark, and bright soliton solutions. Many of these solutions are essential for understanding the behavior of high frequency waves. Such results are tremendously recommended in advanced research and innovation.

REFERENCES

- [1] M. A. Akbar, F. A. Abdullah, and M. M. Khatun, *Optical soliton solutions to the time-fractional Kundu-Eckhaus equation through the $(G'/G, 1/G)$ -expansion technique*, Optical and Quantum Electronics, 55(4) (2023), 291.
- [2] M. A. Akbar, N. H. M. Ali, and J. Hussain, *Optical soliton solutions to the $(2 + 1)$ -dimensional Chaffee–Infante equation and the dimensionless form of the Zakharov equation*, Advances in Difference Equations, 2019(1) (2019), 446.
- [3] T. Aktosun and M. Unlu, *A generalized method for the darbox transformation*, Journal of Mathematical Physics, 63 (2022), 103501.
- [4] A. Ali, J. Ahmad, and S. Javed, *Dynamic investigation to the generalized yu–toda–sasa–fukuyama equation using darbox transformation*, Optical and Quantum Electronics, 56(2) (2024), 166.



- [5] M. Ali, M. A. Khattab, and S. Mabrouk, *Travelling wave solution for the Landau-Ginburg-Higgs model via the inverse scattering transformation method*, Nonlinear Dynamics, *111* (2023), 7687–7697.
- [6] M. R. Ali, M. A. Khattab, and S. Mabrouk, *Investigation of travelling wave solutions for the $(3+1)$ -dimensional hyperbolic nonlinear Schrödinger equation using Riccati equation and F-expansion techniques*, Optical and Quantum Electronics, *55*(11) (2023), 991.
- [7] S. Ali, A. Ullah, S. F. Aldosary, S. Ahmad, and S. Ahmad, *Construction of optical solitary wave solutions and their propagation for Kuralay system using tanh-coth and energy balance method*, Results in Physics, *59* (2024), 107556.
- [8] J. V. Armitage and W. F. Eberlein, *Elliptic functions*, Cambridge University Press, Cambridge, (2006).
- [9] S. Arshed, G. Akram, M. Sadaf, M. Bilal Riaz, and A. Wojciechowski, *Solitary wave behavior of $(2+1)$ -dimensional Chaffee-Infante equation*, Plos one, *18*(1) (2023), e0276961.
- [10] S. Azami and M. Jafari, *Ricci solitons and ricci bi-conformal vector fields on the Lie group $H^2 \times R$* , Reports on Mathematical Physics, *93*(2) (2024), 231-239.
- [11] S. Azami, M. Jafari, A. Haseeb, and A. A. H. Ahmadini, *Cross curvature solitons of Lorentzian three-dimensional Lie groups*, Axioms, *13* (2024), 13040211.
- [12] J. A. Haider, A. M. Alhuthali, and M. A. Elkotb, *Exploring novel applications of stochastic differential equations: Unraveling dynamics in plasma physics with the tanh-coth method*, Results in Physics, *60* (2024), 107684.
- [13] M. Jafari and R. Darvazebanzade, *Analyzing of approximate symmetry and new conservation laws of perturbed generalized Benjamin-Bona-Mahony equation*, AUT Journal of Mathematics and Computing, *5*(1) (2024), 61-69.
- [14] M. Jafari and S. Mahdion, *Analysis of generalized quasilinear hyperbolic and Boussinesq equations from the point of view of potential symmetry*, Journal of Finsler Geometry and its Applications, *5*(1) (2024), 1-10.
- [15] M. Jafari, *New exact non-reducible solutions for generalized Zakharov-Kuznetsov equation*, Journal of Differential Geometry, Applications and its Aspects, *1*(1) (2024), 12-17.
- [16] M. M. A. Khater and B. Ghanbari, *On the solitary wave solutions and physical characterization of gas diffusion in a homogeneous medium via some efficient techniques*, The European Physical Journal Plus, *136*(4) (2021), 447.
- [17] S. Kumar, H. Almusawa, I. Hamid, M. A. Akbar, and M. A. Abdou, *Abundant analytical soliton solutions and evolutionary behaviors of various wave profiles to the chaffee–infante equation with gas diffusion in a homogeneous medium*, Results in Physics, *30* (2021), 104866.
- [18] S. Kumar, H. Almusawa, I. Hamid, and M. Abdou, *Abundant closed-form solutions and solitonic structures to an integrable fifth-order generalized nonlinear evolution equation in plasma physics*, Results in Physics, *26* (2021), 104453.
- [19] S. Kumar, M. Niwas, M. Osman, and M. Abdou, *Abundant different types of exact soliton solution to the $(4+1)$ -dimensional Fokas and $(2+1)$ -dimensional breaking soliton equations*, Communications in Theoretical Physics, *73*(10) (2021), 105007.
- [20] S. Kumar, L. Kaur, and M. Niwas, *Some exact invariant solutions and dynamical structures of multiple solitons for the $(2+1)$ -dimensional bogoyavlensky-konopelchenko equation with variable coefficients using Lie symmetry analysis*, Chinese Journal of Physics, *71* (2021), 518-538.
- [21] Y. Yıldırım and M. Mirzazadeh, *Optical pulses with Kundu-Mukherjee-Naskar model in fiber communication systems*, Chinese Journal of Physics, *64* (2020), 183-193.
- [22] C. Li, L. Chen, and G. Li, *Optical solitons of space-time fractional Sasa-Satsuma equation by F-expansion method*, Optik, *224* (2020), 165527.
- [23] S. Mabrouk and A. Rashed, *On the G'/G expansion method applied to $(2+1)$ -dimensional asymmetric Nizhnik-Novikov-Veselov equation*, Journal of Advances in Applied & Computational Mathematics, *10* (2023), 39-49.
- [24] S. Mabrouk, A. Rashed, and R. Saleh, *Unveiling traveling waves and solitons of Dirac integrable system via homogenous balance and singular manifolds methods*, Computational Methods for Differential Equations, (2024).
- [25] S. Mabrouk, M. Mahdy, A. Rashed, and R. Saleh, *Exploring high-frequency waves and soliton solutions of fluid turbulence through relaxation medium modeled by Vakhnenko-Parkes equation*, Computational Methods for Differential Equations, (2024).



- [26] A. Mahmood, M. Abbas, G. Akram, M. Sadaf, M. B. Riaz, and T. Abdeljawad, *Solitary wave solution of (2+1)-dimensional Chaffee-Infante equation using the modified Khater method*, Results in Physics, 48 (2023), 106416.
- [27] M. Mohamed, S. M. Mabrouk, and A. S. Rashed, *Mathematical investigation of the infection dynamics of covid-19 using the fractional differential quadrature method*, Computation, 11 (10) (2023), 198.
- [28] A. S. Rashed, *Interaction of two long waves in shallow water using hirota-satsuma model and similarity transformations*, Delta University Scientific Journal, 5(1) (2022), 93-101.
- [29] A. S. Rashed, M. Inc, and R. Saleh, *Extensive novel waves evolution of three-dimensional Yu-Toda-Sasa-Fukuyama equation compatible with plasma and electromagnetic applications*, Modern Physics Letters B, 37(1) (2023), 2250195.
- [30] A. S. Rashed, M. Mohamed, and M. Inc, *Plasma particles dispersion based on Bogoyavlensky-Konopelchenko mathematical model*, Computational Methods for Differential Equations, 12(3) (2024), 561-570.
- [31] A. S. Rashed, S. M. Mabrouk, and A. M. Wazwaz, *Forward scattering for non-linear wave propagation in (3 + 1)-dimensional jimbo-miwa equation using singular manifold and group transformation methods*, Waves in Random and Complex Media, 32(2) (2022), 663-675.
- [32] M. Sadaf, S. Arshed, and G. Akram, *Exact soliton and solitary wave solutions to the fokas system using two variables ($G'/G, 1/G$)-expansion technique and generalized projective Riccati equation method*, Optik, 268 (2022), 169713.
- [33] R. Saleh and A. S. Rashed, *New exact solutions of (3 + 1)-dimensional generalized Kadomtsevpetviashvili equation using a combination of Lie symmetry and singular manifold methods*, Mathematical Methods in the Applied Sciences, 43(4) (2020), 2045-2055.
- [34] R. Saleh, A. S. Rashed, and A. M. Wazwaz, *Plasma-waves evolution and propagation modeled by sixth order Ramani and coupled Ramani equations using symmetry methods*, Physica Scripta, 96(8) (2021), 085213.
- [35] Y. Wang and X. Lü, *Bäcklund transformation and interaction solutions of a generalized Kadomtsev-Petviashvili equation with variable coefficients*, Chinese Journal of Physics, 89 (2024), 37-45.
- [36] S. Kumar, W. X. Ma, and A. Kumar, *Lie symmetries, optimal system and group-invariant solutions of the (3+1)-dimensional generalized KP equation*, Chinese Journal of Physics, 69 (2021), 1-23.
- [37] Y. Yıldırım, A. Biswas, M. Ekici, E. M. E. Zayed, S. Khan, L. Moraru, A. K. Alzahrani, and M. R. Belic, *Highly dispersive optical solitons in birefringent fibers with four forms of nonlinear refractive index by three prolific integration schemes*, Optik, 220 (2020), 165039.
- [38] Y. Yıldırım, *Optical solitons with Biswas-Arshed equation by F-expansion method*, Optik, 227 (2021), 165788.
- [39] A. Yusuf, T. A. Sulaiman, A. Abdeljabbar, and M. Alquran, *Breather waves, analytical solutions and conservation laws using Lie-Bäcklund symmetries to the (2+1)-dimensional Chaffee-Infante equation*, Journal of Ocean Engineering and Science, 8(2) (2023), 145-151.
- [40] L. Zhang, C. Li, and H. Wang, *Bäcklund transformations of multi-component Boussinesq and Degasperis-Procesi equations*, International Journal of Geometric Methods in Modern Physics, 21(3) (2024), 2450066.
- [41] Y. Zhou, M. Wang, and Y. Wang, *Periodic wave solutions to a coupled KdV equations with variable coefficients*, Physics Letters A, 308(1) (2003), 31-36.

